

Differential Geometry of Curves and Surfaces in Lorentz-Minkowski space

Mini-Course taught at the
Instituto de Matemática e Estatística (IME-USP)
University of Sao Paulo, Brasil

Rafael López

Departamento de Geometría y Topología
Universidad de Granada
18071 Granada, España
e-mail: rcamino@ugr.es

October 18, 2008

Contents

Preface	i
1 The Lorentz-Minkowski space $\mathbf{E}_1^3$	1
1.1 Basic definitions	1
1.2 Timelike vectors	6
1.3 The Lorentzian vector product	9
1.4 Isometries of $\mathbf{E}_1^3$	10
1.5 Exercises	13
2 Curves in Minkowski space	17
2.1 Parametrized curves	17
2.2 Curvature and torsion	20
2.2.1 The timelike case	21
2.2.2 The spacelike case	21
2.2.3 The lightlike case	22
2.3 Planar curves with constant curvature	23
2.4 Helices and Bertrand curves in $\mathbf{E}_1^3$	25
2.5 Exercises	28
3 Spacelike and timelike surfaces in $\mathbf{E}_1^3$	29
3.1 Surfaces in $\mathbf{E}_1^3$	29
3.2 Mean curvature of non-degenerate surfaces	32
3.3 Local computations of curvatures	35
3.4 Umbilical surfaces	36
3.5 Exercises	38

4	Spacelike surfaces with constant mean curvature	39
4.1	The variational problem	39
4.2	The maximum principle	42
4.3	Two equations for CMC spacelike surfaces	44
5	Compact surfaces with constant mean curvature	47
5.1	Consequences of the maximum principle	47
5.2	The Dirichlet problem: the Euclidean case	50
5.3	The Dirichlet problem: the Lorentzian case	53
5.4	Exercises	54
6	Lorentzian Riemann examples	57
6.1	Introduction to the problem	57
6.2	The planes of the foliation are parallel	59
6.3	The planes of the foliation are not parallel	62
6.4	Exercises	64
	Bibliography	65

Preface

This set of notes is a written version of a Mini-Course given in September 2008 while I was visiting for two weeks the Instituto de Matemática e Estatística (IME-USP) of the Sao Paulo University, Brasil. The course of eight hours was designed for undergraduate and graduate students in Mathematics. Background knowledge of differential geometry of curves and surfaces will be assumed, basically as in Do Carmo's textbook.

This Mini-Course gives an introduction to classical differential geometry of curves and surfaces in Lorentz-Minkowski space $\mathbf{E}_1^3$. In the case of surfaces, we will study spacelike surfaces, specially with the assumption that its mean curvature is constant. Throughout the lectures, we will compare the results and techniques with those ones in Euclidean ambient space.

The topics that this Mini-Course covers will be self-contained, in such way the each chapter tries to use the material of the previous ones. We begin in Chapter 1 with an introduction to the metric space of Lorentz-Minkowski space $\mathbf{E}_1^3$ with some details on the isometries of this space. The second chapter is devoted to develop the Frenet equations for curves in $\mathbf{E}_1^3$. This part follows the Euclidean notions, such as, planar curves with constant curvature, helices and Bertrand curves.

In Chapter 3 we begin with the study of surfaces in $\mathbf{E}_1^3$ and we pay special attention on spacelike surfaces. We give the notion of mean curvature and we find all umbilical surfaces in $\mathbf{E}_1^3$. In Chapter 4 we characterize spacelike surfaces with constant mean curvature as solutions of a variational problem. Moreover, we give the maximum principle for this kind of surfaces, which will be used in Chapter 5 to study the shape of a spacelike compact surface with constant mean curvature. We give the basic tools to solve the Dirichlet problem and we compare the solvability with what happens in Euclidean space. Finally, Chapter 6 is devoted to study spacelike surfaces with constant mean curvature and foliated by circles. We show the Lorentzian catenoid as well as the corresponding Riemann examples for the Lorentzian space.

A final remark. The reader can find lacks in the topics that cover these lectures. However, our plan was to write the Mini-Course and we do not expect to realize a (definitive) text of differential geometry in Minkowski space. Also, he will find typographic errors and an English wording not so polished. Anyway, the author of this text will thank any suggestion to improve this version (e-mail address: rcamino@ugr.es). There is a Spanish version of these notes upon request.

Chapter 1

The Lorentz-Minkowski space $\mathbf{E}_1^3$

1.1 Basic definitions

Let $\mathbb{R}^3$ be the real vector space with its usual vector structure. Denote by $B = \{E_1, E_2, E_3\}$ the usual base of $\mathbb{R}^3$, that is,

$$E_1 = (1, 0, 0), \quad E_2 = (0, 1, 0), \quad E_3 = (0, 0, 1).$$

We will use (x, y, z) or (x_1, x_2, x_3) to denote the coordinates of a vector with respect to B . We also consider in $\mathbb{R}^3$ its affine structure, and we will say "horizontal" or "vertical" in its usual sense.

If $\{e_1, \dots, e_m\}$ is a finite set of vector, we denote by $\langle e_1, \dots, e_m \rangle$ the vector subspace spanning by their linear combinations, that is, $\langle e_1, \dots, e_m \rangle = \{\sum a_i e_i; a_i \in \mathbb{R}, 1 \leq i \leq m\}$.

Definition 1.1.1. *The Lorentz-Minkowski space is the metric space $\mathbf{E}_1^3 = (\mathbb{R}^3, \langle, \rangle)$, where the metric $\langle, \rangle$ is given by*

$$\langle u, v \rangle = u_1 v_1 + u_2 v_2 - u_3 v_3, \quad u = (u_1, u_2, u_3), v = (v_1, v_2, v_3).$$

The metric $\langle, \rangle$ is called the Lorentzian metric¹.

We remark that $\langle, \rangle$ is a non-degenerate metric of index 1. We also call $\mathbf{E}_1^3$ as Minkowski space, and $\langle, \rangle$ as the Minkowski metric. We also write

$$\langle u, v \rangle = u^t \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} v := u^t G v.$$

The vector space $\mathbb{R}^3$ also supports the Euclidean metric, which will be denoted in this text by $\langle, \rangle_\epsilon$. We write $\mathbf{E}^3$ the metric space $(\mathbb{R}^3, \langle, \rangle_\epsilon)$ to distinguish from the Lorentz-Minkowski space².

¹In the literature, Minkowski space is sometimes denoted by $\mathbb{L}^3$. Also it appears the metric defined as $\langle(u_0, u_1, u_2), (v_0, v_1, v_2)\rangle = -u_0 v_0 + u_1 v_1 + u_2 v_2$, and the index 0 means the place where the minus sign lies in the metric

²Now, the index 1 in the symbol $\mathbf{E}_1^3$ has its sense by meaning that the metric has index 1

A great number of the next definitions and results can generalize in higher dimensions, that is, to the space $\mathbf{E}_1^n = (\mathbb{R}^n, \langle, \rangle)$, with the usual Lorentzian metric.

If we consider an orthonormal base $B = \{e_1, e_2, e_3\}$, that is, a base where the metric diagonalizes with 1 and -1 , we always arrange in such way that the matricial expression of the metric is $\text{diag}[1, 1 - 1]$. In general, for a given base B , we denote the metric coefficients by $g_{ij} = \langle e_i, e_j \rangle$.

Definition 1.1.2. A vector $v \in \mathbf{E}_1^3$ is called

1. spacelike if $\langle v, v \rangle > 0$ or $v = 0$,
2. timelike if $\langle v, v \rangle < 0$ and
3. lightlike if $\langle v, v \rangle = 0$ and $v \neq 0$.

We point out that the null vector $v = 0$ is considered of spacelike type although it satisfies $\langle v, v \rangle = 0$. The *light-cone* of $\mathbf{E}_1^3$ is defined as the set of all lightlike vectors of $\mathbf{E}_1^3$, that is,

$$\mathcal{C} = \{(x, y, z) \in \mathbf{E}_1^3; x^2 + y^2 - z^2 = 0\} - \{(0, 0, 0)\}.$$

See figure 1.1. The set of timelike vectors will be denote by $\mathcal{T}$ and it is the following set:

$$\mathcal{T} = \{(x, y, z) \in \mathbf{E}_1^3; x^2 + y^2 - z^2 < 0\}.$$




Figure 1.1: The causal character in Minkowski space.

Given $U \subset \mathbb{R}^3$ a vector subspace, we consider on U the induced metric $\langle, \rangle_U$:

$$\langle u, v \rangle_U = \langle u, v \rangle, \quad u, v \in U.$$

From now, we drop the subindex U . The subspace U is called spacelike (resp. timelike or lightlike) if the induced metric is positive definite (resp. non-degenerate of index 1, degenerate and $U \neq \{0\}$).

The *causal character* of a vector or a subspace is the property to be spacelike, timelike or lightlike. Any subspace belongs to one of the above three kinds.

Examples.

1. The vectors E_1 and E_2 are spacelike; the vector E_3 is timelike; the vector $E_2 + E_3$ is lightlike.
2. The plane $\langle E_1, E_2 \rangle$ is spacelike; the planes $\langle E_1, E_3 \rangle$ and $\langle E_2, E_3 \rangle$ are timelike; the plane $\langle E_1, E_2 + E_3 \rangle$ is lightlike.
3. The vector $E_1 + E_2 + E_3$ is spacelike but the plane $\langle E_1, E_1 + E_2 + E_3 \rangle$ is lightlike.
4. The vector $E_2 + E_3$ is lightlike, but the plane $\langle E_2 + E_3, E_3 \rangle$ is timelike.

Let us recall some facts that happen in a metric space (V, g) with a non-degenerate metric. As usually, $U^\perp$ denotes the orthogonal subspace to U , that is, $U^\perp = \{v \in V, g(u, v) = 0, \forall u \in U\}$.

Lemma 1.1.3. *Let (V, g) be a metric space where g is a non-degenerate metric.*

1. *If $U \subset V$ is a subspace, then $\dim(U^\perp) = \dim(V) - \dim(U)$.*
2. *If $U \subset V$ is a subspace, then $(U^\perp)^\perp = U$.*
3. *If $U \subset V$ is a non-degenerate subspace, then $U^\perp$ is a non-degenerate subspace too.*

Proof:

1. Let $\{e_1, \dots, e_m\}$ a base of U and we extend it until to be a base $B = \{e_1, \dots, e_n\}$ of V . If $u = \sum_i x_i e_i \in U^\perp$, then

$$0 = \left\langle \sum_{i=1}^n x_i e_i, e_j \right\rangle = \sum_{i=1}^n g_{ji} x_i = 0, \quad 1 \leq j \leq m.$$

In a matricial expression, these m -equations write as

$$\begin{pmatrix} g_{11} & \dots & g_{1n} \\ \dots & \dots & \dots \\ g_{m1} & \dots & g_{mn} \end{pmatrix} \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}$$

or $AX = 0$, and $A = (g_{ij})_{m \times n}$. The range of A is m because there is a sub-matrix with range exactly n (this is due to the non-degeneracy of the metric). As consequence, the solutions of $AX = 0$ generate a $n - m$ -dimensional subspace.

2. Since $(U^\perp)^\perp \subset U$, the result follows from $\dim(U^\perp)^\perp = \dim(U)$.
3. Let $B = \{e_1, \dots, e_m\}$ be an orthonormal base of U , that is, the matrix of the metric $g|_U$ is diagonal with only 1 and -1 . We extend the base to get an orthonormal base of V , namely, $B = \{e_1, \dots, e_n\}$. Since $\dim(U^\perp) = n - m$, then $\{e_{m+1}, \dots, e_n\}$ is a base of $U^\perp$ and this ends the proof.

□

We now give characterizations of subspaces depending on its causal character.

Proposition 1.1.4. *1. Let $v \in \mathbf{E}_1^3$. Then v is a timelike vector if and only if $\langle v \rangle^\perp$ is spacelike and so, $\mathbf{E}_1^3 = \langle v \rangle \oplus \langle v \rangle^\perp$. For spacelike vectors, we have: v is spacelike if and only if $\langle v \rangle^\perp$ is timelike.*

2. Let $U \subset V$ be a subspace. Then U is spacelike if and only if $U^\perp$ is timelike.
3. If U is a subspace, then U is lightlike if and only if $U^\perp$ is lightlike too.

Proof:

1. If v is a timelike vector, and by multiplying by a number if it is necessary, we put v as a part of an orthonormal base of $\mathbf{E}_1^3$, $B = \{e_1, e_2, v\}$. Then $\langle v \rangle^\perp = \langle e_1, e_2 \rangle$, which it is a spacelike subspace. For the converse, let $\{e_1, e_2\}$ be a orthonormal base of $\langle v \rangle^\perp$, where $\langle \cdot, \cdot \rangle_{|\langle v \rangle^\perp}$ is a positive definite metric. Then $\{e_1, e_2, v\}$ is a base where diagonalizes the metric. As $g_{11} = g_{22} = 1$, then $g_{33} < 0$, that is, v is a timelike vector.
2. If U is a timelike subspace, let $v \in U$ be a timelike vector. Then $U^\perp \subset \langle v \rangle^\perp$. As $\langle v \rangle^\perp$ is spacelike, then $U^\perp$ is spacelike. The converse is analogous, together the fact that $(U^\perp)^\perp = U$.
3. This item is a consequence of the above two ones.

□

For the usual sense that one has of Euclidean space $\mathbf{E}^3$, Lorentz-Minkowski space presents us as some "strange" due to the existence of timelike vectors and mainly, by the fact of lightlike vectors, that is, vectors whose product by itself vanishes³. We now study properties of lightlike vectors.

- Proposition 1.1.5.** *1. If u and v are two lightlike vectors, then they are dependent linear and only $\langle u, v \rangle = 0$.*
- 2. If u and v are two timelike or lightlike vectors with $\langle u, v \rangle = 0$, then they are lightlike vectors.*
- 3. If U is a lightlike subspace, then $\dim(U \cap U^\perp) = 1$.*

Proof:

1. If u and v are proportional, then they are orthogonal. Now, we suppose that they are orthogonal. In the decomposition $\mathbf{E}_1^3 = \langle E_3 \rangle^\perp \oplus \langle E_3 \rangle$, we write $u = x + w$ and $v = y + w$: we assume that the vector w is the same in both decomposition in order what to want to show. As $\langle u, v \rangle = 0$ and both are lightlike vectors, then

$$\langle x, y \rangle + \langle w, w \rangle + \langle x, w \rangle + \langle y, w \rangle = 0.$$

$$\langle x, x \rangle + \langle w, w \rangle + 2\langle x, w \rangle = 0.$$

$$\langle y, y \rangle + \langle w, w \rangle + 2\langle y, w \rangle = 0.$$

By combining the three equations, we obtain $|x|^2 + |y|^2 - 2\langle x, y \rangle = 0$, that is, $|x - y|^2 = 0$. Thus $x = y$, because $x - y$ is a spacelike vector ($x - y \in \langle w \rangle^\perp$). So we deduce $u = v$.

2. If the two vectors are timelike, then $\langle u, v \rangle \neq 0$. This is due to the following fact: by using $\mathbf{E}_1^3 = \langle v \rangle^\perp \oplus \langle v \rangle$, where $\langle v \rangle^\perp$ is a spacelike subspace, we write $u = x + \lambda v$; then

$$\langle u, v \rangle = \langle v, x \rangle + \lambda \langle v, v \rangle = \lambda \langle v, v \rangle.$$

If $\langle u, v \rangle = 0$, then $\lambda = 0$ and $u = x$ would be spacelike. The reasoning is analogous with lightlike or timelike vectors. Therefore both vectors must be lightlike.

³This difficulty is similar, in a some sense, with what happens for the set of numbers: there exists complex numbers x such that $x^2 = -1$, but there are not complex numbers, $x \neq 0$, such that $x^2 = 0$.

3. If $u, v \in U \cap U^\perp$, then $\langle u, v \rangle = 0$. Thus they are dependent linear. This proves that $\dim(U \cap U^\perp) \leq 1$. If the dimension is exactly 0, then $\mathbf{E}_1^3 = U \oplus U^\perp$, and so any vector of $\mathbf{E}_1^3$ would be lightlike.

□

We study timelike subspaces.

Proposition 1.1.6. *Let $U \subset \mathbf{E}_1^3$ be a two-dimensional subspace. The following statements are equivalent:*

1. U is a timelike subspace.
2. U contains two independent linear lightlike vectors.
3. U contains a timelike vector.

Proof:

- 1 $\Rightarrow$ 2 Let $\{e_1, e_2, e_3\}$ be an orthonormal base of $\mathbf{E}_1^3$. Then $e_2 + e_3$ and $e_2 - e_3$ are independent linear, lightlike vectors.
- 2 $\Rightarrow$ 3 If u and v are the two independent linear, lightlike vectors, then $u + v$ or $u - v$ is a timelike vector because
- $$\langle u \pm v, u \pm v \rangle = \pm 2\langle u, v \rangle$$
- and $\langle u, v \rangle \neq 0$ due to both vectors are timelike.
- 3 $\Rightarrow$ 1 Let v be a timelike vector of U . Then $U^\perp \subset \langle v \rangle^\perp$, and $\langle v \rangle^\perp$ is a spacelike subspace. Thus, $U^\perp$ is spacelike, and so, U is timelike.

□

The above result can generalize to arbitrary dimensions by considering that U is a hyperplane. We now characterize lightlike subspaces.

Proposition 1.1.7. *Let U be a subspace of $\mathbf{E}_1^3$. The following statements are equivalent:*

1. U is a lightlike subspace.
2. U contains a lightlike vector but not a timelike one.
3. $U \cap \mathcal{C} = L - \{0\}$, and $\dim L = 1$.

Proof:

- 1 $\Rightarrow$ 2 Since $\langle, \rangle$ is a degenerate metric, there is a lightlike vector. By the Proposition 1.1.6, there are not timelike vectors.
- 2 $\Rightarrow$ 3 Since there exist lightlike vectors, $U \cap \mathcal{C}$ is a non-empty set. By using Proposition 1.1.6 again, if there are two independent linear, lightlike vectors, then there would be a timelike vector.

3 $\Rightarrow$ 1 Proposition 1.1.6 says that U is not a spacelike nor timelike subspace. □

Proposition 1.1.8. *Let P be a plane of $\mathbf{E}_1^3$. We denote by $\vec{n}$ an orthogonal vector with the Euclidean metric. Then P is a spacelike (resp. timelike, lightlike) plane if and only if $\vec{n}$ is a timelike (resp. spacelike, lightlike) vector.*

Proof: If P writes as $P = \{(x, y, z) \in \mathbb{R}^3; ax + by + cz = 0\}$, then $\vec{n}$ is proportional to the vector (a, b, c) . We can also write P as

$$P = \{(x, y, z) \in \mathbb{R}^3; ax + by - (-c)z = 0\} = \langle (a, b, -c) \rangle^\perp.$$

The causal character of $(a, b, -c)$ is the same than $\vec{n}$, which proves the result. □

We give the definition of the modulus of a vector.

Definition 1.1.9. *Given $u \in \mathbf{E}_1^3$, we call the modulus or norm of u as the number $\sqrt{|\langle u, u \rangle|}$ and we write $|u|$. The vector u is called unitary if its modulus is 1.*

As a consequence, if u is a spacelike vector (resp. timelike), we have $|u| = \sqrt{\langle u, u \rangle}$ (resp. $|u| = \sqrt{-\langle u, u \rangle}$).

Proposition 1.1.10. *If P is a spacelike plane and $P = \langle v \rangle^\perp$, with $\langle v, v \rangle = -1$, we have*

$$|v|_\epsilon \geq 1,$$

where the subindex ϵ represents that the computations are done with the Euclidean metric of $\mathbb{R}^3$.

Proof: We write $P = \{(x, y, z) \in \mathbb{R}^3; ax + by + cz = 0\}$, with $\vec{n} = (a, b, c)$ and $a^2 + b^2 + c^2 = 1$. Then $P = \langle v \rangle^\perp$, where

$$v = \frac{(a, b, -c)}{\sqrt{c^2 - a^2 - b^2}}$$

and it satisfies $\langle v, v \rangle = -1$. By computing the Euclidean norm of v , we obtain

$$|v|_\epsilon^2 = \frac{a^2 + b^2 + c^2}{c^2 - a^2 - b^2} = \frac{1}{c^2 - a^2 - b^2} \geq 1.$$

□

This result justifies why when one draws the orthogonal vector to a spacelike plane, the (Euclidean) size is greater than the Euclidean unit orthogonal vector. See figure 1.2

1.2 Timelike vectors

Let us recall that $\mathcal{T}$ is the set of timelike vectors of $\mathbf{E}_1^3$. For each $u \in \mathcal{T}$, we define the *timelike cone* of u as the set given by

$$C(u) = \{v \in \mathcal{T}; \langle u, v \rangle < 0\}.$$

This set is non-empty since $u \in C(u)$. Moreover $\mathcal{T}$ is the disjoint union of $C(u)$ and $C(-u)$: if $v \in \mathcal{T}$, then $\langle u, v \rangle \neq 0$, and so $v \in C(u)$ or $v \in C(-u)$. Furthermore $C(u) \cap C(-u) = \emptyset$. Some properties of timelike cones are:




Figure 1.2: The vector v orthogonal to a spacelike plane P "appears" bigger than the Euclidean normal vector $\vec{n}$ to P

- Proposition 1.2.1.**
1. Two timelike vectors u and v lie in the same timelike cone if and only if $\langle u, v \rangle < 0$.
 2. $u \in C(v)$ if and only if $C(u) = C(v)$.
 3. The timelike cones are convex sets.

Proof:

1. If $\langle u, v \rangle < 0$, then $u \in C(v)$. Let assume that $u, v \in C(w)$. We can suppose that $\langle w, w \rangle = -1$. We write $u = x + aw$ and $v = y + bw$, with $x, y \in w^\perp$. As $w^\perp$ is a spacelike subspace, then $|\langle x, y \rangle| \leq |x||y|$, and

$$\langle u, v \rangle = -ab + \langle x, y \rangle \leq -ab + |x||y|.$$

But $\langle x, x \rangle < a^2$ and $\langle y, y \rangle < b^2$.

2. If $u \in C(v)$, then $\langle u, v \rangle < 0$, that is $v \in C(u)$.
3. Assume that $u, v \in C(w)$ and let $t \in [0, 1]$. Then

$$\langle tu + (1-t)v, w \rangle = t\langle u, w \rangle + (1-t)\langle v, w \rangle < 0,$$

and this means that $tu + (1-t)v \in C(w)$.

□

We now show a type of Cauchy-Schwarz inequality for timelike vectors. This inequality allows us to define the angle between two timelike vectors.

Theorem 1.2.2. Let u and v two timelike vectors. Then

$$|\langle u, v \rangle| \geq \sqrt{-\langle u, u \rangle} \sqrt{-\langle v, v \rangle}$$

and the equality holds if and only if u and v are two proportional vectors. In the case that both vectors lie in the same timelike cone, there exists a unique number $\varphi \geq 0$ such that

$$\langle u, v \rangle = -|u||v| \cosh \varphi.$$

This number φ is called the hyperbolic angle between u and v .

Proof: We consider two independent timelike vectors u and v . Then $U = \langle u, v \rangle$ is a timelike plane. By Proposition 1.1.6 the equation on a and b given by

$$\langle au + bv, au + bv \rangle = a^2 \langle u, u \rangle + b^2 \langle v, v \rangle + 2ab \langle u, v \rangle = 0$$

has solution. Moreover $a \neq 0$ (on the contrary, the vector v would be timelike). By dividing by a , we have that

$$\langle u, u \rangle + 2\lambda \langle u, v \rangle + \lambda^2 \langle v, v \rangle = 0$$

has solution on λ . In particular, the discriminant of the quadratic equation must be positive, that is,

$$\langle u, v \rangle^2 > \langle u, u \rangle \langle v, v \rangle.$$

This shows the inequality in the case that u and v are independent linear. On the other hand, if they are proportional, then we obtain equality.

For the second part of the theorem, we write

$$\frac{\langle u, v \rangle^2}{(-\langle u, u \rangle)(-\langle v, v \rangle)} \geq 1. \quad (1.1)$$

If u and v lie in the same timelike cone, then $\langle u, v \rangle < 0$ and the expression (1.1) implies

$$\frac{-\langle u, v \rangle}{\sqrt{-\langle u, u \rangle} \sqrt{-\langle v, v \rangle}} \geq 1.$$

As the hyperbolic cosine function $\cosh : [0, \infty) \rightarrow [1, \infty)$ is one-to-one, there exists a unique number φ such that

$$\cosh \varphi = \frac{-\langle u, v \rangle}{\sqrt{-\langle u, u \rangle} \sqrt{-\langle v, v \rangle}}.$$

□

Corollary 1.2.3. *If u, v are two timelike vectors that lie in the timelike cone, then*

$$|u + v| \geq |u| + |v|$$

and the equality holds if and only if u and v are proportional.

Let us see that in general there is not a notion of angle for spacelike vectors. For example, for the unit vectors $u = (0, \cosh(t), \sinh(t))$ and $v = (0, 1, 0)$, we have $\langle u, v \rangle = \cosh(t)$, which attains any arbitrary value (and greater than 1). This is due to the plane $\langle u, v \rangle$ is timelike. On the contrary, if u and v determine a spacelike plane, the induced metric on P is positive definite and then, we would have the (usual) Cauchy-Schwarz inequality.

We end this section with the definition of *timelike orientation*. Before, we recall the notion of orientation in any vector space. For this, in the set of all bases of $\mathbb{R}^3$, we consider the equivalence relation R given by BRB' if the matrix that changes both bases has positive determinant. There exist exactly two equivalence classes, called orientations of $\mathbb{R}^3$. When we fix one of them, we say that $\mathbb{R}^3$ is oriented (with this orientation). Exactly, when we say that $\mathbb{R}^3$ is oriented we mean the ordered pair $(\mathbb{R}^3, [B])$, where B is a base. In such case, if B' is any base, we say that B' is positive oriented if $B' \in [B]$; on the contrary, we say that B' is negative oriented.

In Minkowski space $\mathbf{E}_1^3$ there is not sense to talk again of orientation, since this is defined in $\mathbb{R}^3$ as vector space, but not as metric space. The timelike orientation that we want to introduce is a metric concept, because we use the Lorentzian metric $\langle \cdot, \cdot \rangle$. Thus, there is not relation between both notions.

In $\mathbf{E}_1^3$ we consider the set β of all orthonormal bases. We define the equivalence relation given by

$$B \sim B' \text{ if } e_3 \text{ and } e'_3 \text{ lies in the same timelike cone,}$$

that is, if $\langle e_3, e'_3 \rangle < 0$. We have only two equivalence classes, which are called *timelike orientations*. Moreover, each class determines a unique timelike cone and conversely, given a timelike cone, there exists a unique timelike orientation in such way that all bases belonging this orientation have the last vector in such timelike cone.

We say that $\mathbf{E}_1^3$ is *timelike oriented* if we fix an orientation, that is, we consider the ordered pair $(\mathbf{E}_1^3, [B])$, for some B .

Definition 1.2.4. Let $E_3 = (0, 0, 1)$. Given a timelike vector v , we say that v is *future-directed* (resp. *past-directed*) if $v \in C(E_3)$, that is, $\langle v, E_3 \rangle < 0$ (resp. $v \in C(-E_3)$, or $\langle v, E_3 \rangle > 0$).

It is also equivalent to say that $v = (v_1, v_2, v_3)$ is future-directed if $v_3 > 0$. We always orient $\mathbf{E}_1^3$ by the timelike cone $C(E_3)$, that is, $(\mathbf{E}_1^3, [B_u])$, where B_u is the usual base of $\mathbb{R}^3$.

1.3 The Lorentzian vector product

The definition of vector product is the same that the given one in the Euclidean ambient.

Definition 1.3.1. If $u, v \in \mathbf{E}_1^3$, we call the (Lorentzian) vector product of u and v to the unique vector denoted by $u \times v$ that satisfies

$$\langle u \times v, w \rangle = \det(u, v, w), \quad (1.2)$$

where $\det(u, v, w)$ is the determinant of the matrix obtained by putting by columns the coordinates of the three vectors u , v and w .

By taking w each one of the vectors of the usual base, we obtain

$$u \times v = \begin{vmatrix} i & j & -k \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}. \quad (1.3)$$

The bilinearity of the metric assures the existence and uniqueness of this vector. Thus, if we denote by $u \times_\epsilon v$ the Euclidean vector product, we have that $u \times v$ is the reflection of $u \times_\epsilon v$ with respect to the plane $\{z = 0\}$.

Proposition 1.3.2. *The vector product satisfies the following properties:*

1. $u \times v = -v \times u$.
2. $u \times v$ is orthogonal to u and v .
3. $u \times v = 0$ if and only if $\{u, v\}$ are not proportional.
4. $u \times v \neq 0$ lies in the plane $P = \langle u, v \rangle$ if and only if the plane P is lightlike,

1.4 Isometries of $\mathbf{E}_1^3$

In this section we study the isometries of Minkowski space $\mathbf{E}_1^3$. We consider the set of all vector isometries of $\mathbf{E}_1^3$, and we denote it by $O_1(3)$. If B and B' are two orthonormal bases, the matrix A of change of coordinates satisfies $A^t G A = G$. Thus

$$O_1(3) = \{A \in Gl(3, \mathbb{R}); A^t G A = G\}.$$

In particular, $\det(A) = \pm 1$. This means that $O_1(3)$ has at least two connected components⁴. We denote by $SO_1(3)$ the set of isometries with determinant 1. The set $SO_1(3)$ is called the *special Lorentz group*. This group appears related with the notion of orientation of $\mathbb{R}^3$. Exactly, if we fix the orientation given by the usual base, and if B is an orthonormal base, we have $B \in SO_1(3)$ if and only if B is positive oriented.

We define the *ortocrone group* by

$$O_1^+(3) = \{A \in O_1(3); A \text{ maintains the timelike orientation}\}.$$

We say that A maintains the timelike orientation if given a future-directed orthonormal base B , then the base obtained by $B' = A \cdot B$ is also future-directed. We also have the next characterization of $O_1^+(3)$: $A \in O_1^+(3)$ if and only $a_{33} > 0$. The set $O_1^+(3)$ is a group with two components: one of them is $O_1^+(3) \cap SO_1(3)$ and the other one is $O_1^+(3) - (O_1^+(3) \cap SO_1(3))$.⁵

Our interest is the following group. We define the *special Lorentz ortocrone group* as the set

$$O_1^{++}(3) = SO_1(3) \cap O_1^+(3) = \{A \in O_1(3); \det(A) = 1, A \text{ maintains the timelike orientation}\}.$$

This set is a group and $I \in O_1^{++}(3)$. From a topological viewpoint, $O_1^{++}(3)$ is not a compact set because the subset

$$\left\{ \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cosh(t) & \sinh(t) \\ 0 & \sinh(t) & \cosh(t) \end{pmatrix}; t \in \mathbb{R} \right\}$$

is not bounded.

Without proof, we have the following

⁴We are considering in $Gl(3, \mathbb{R})$ the usual topology, that is, the induced by the Euclidean one from $\mathbf{E}^9$ and the inclusion map $Gl(3, \mathbb{R}) \subset \mathbb{R}^9$

⁵Recall that the isometries of $\mathbf{E}^3$, the orthogonal group $O(3)$, it has exactly two connected components, being one of them, the special orthogonal group $SO(3)$. Moreover $SO(3)$ is a compact set.

Theorem 1.4.1. *The connected components of $O_1(3)$ are $O_1^{++}(3)$ y*

$$\begin{aligned} O_1^{+-}(3) &= \{A \in SO_1(3); a_{33} < 0\} \\ O_1^{-+}(3) &= \{A \in O_1^+(3); \det(A) = -1\} \\ O_1^{--}(3) &= \{A \in O_1(3); \det(A) = -1, a_{33} < 0\} \end{aligned}$$

If we denote by T_1 and T_2 the isometries given by $T_1 = \text{diag } [1, 1, -1]$ and $T_2 = \text{diag } [1, -1, 1]$, then the three last components correspond, respectively, with $T_1 \cdot T_2 \cdot O_1^{++}(3)$, $T_2 \cdot O_1^{++}(3)$ y $T_1 \cdot O_1^{++}(3)$.

The rigid motions of $\mathbf{E}_1^3$ are the composition of a vector isometry and a translations of $\mathbf{E}_1^3$.

Next we study the isometries of the two-dimensional Lorentz-Minkowski space $\mathbf{E}_1^2$. This will clarify why we have to distinguish between isometries that maintain or not the timelike orientation. Let A be a matrix given by $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. Then $A \in O_1(2)$ if and only if $G = A^t G A$, that is,

$$a^2 - c^2 = 1, \quad ab - cd = 0, \quad d^2 - b^2 = 1.$$

From the first equation, we have two possibilities:

1. There exists t such that $a = \cosh(t)$ and $c = \sinh(t)$. From $d^2 - b^2 = 1$, it appears two cases again:
 - (a) There exists s such that $d = \cosh(s)$ and $b = \sinh(s)$. With the second equation, we conclude that $s = t$.
 - (b) There exists s such that $d = -\cosh(s)$ y $b = \sinh(s)$. Now we have $s = -t$.
2. There exists t such that $a = -\cosh(t)$ and $c = \sinh(t)$. Equation $d^2 - b^2 = 1$ yields two possibilities:
 - (a) There exists s such that $d = \cosh(s)$ and $b = \sinh(s)$. The second equation concludes that $s = -t$.
 - (b) There exists s such that $d = -\cosh(s)$ and $b = \sinh(s)$. From $ab - cd = 0$, we have $s = t$.

As conclusion, we obtain four kinds of isometries. In the same order as above, they are the following:

$$\begin{aligned} &\begin{pmatrix} \cosh(t) & \sinh(t) \\ \sinh(t) & \cosh(t) \end{pmatrix}, \begin{pmatrix} \cosh(t) & \sinh(t) \\ -\sinh(t) & -\cosh(t) \end{pmatrix}, \\ &\begin{pmatrix} -\cosh(t) & \sinh(t) \\ -\sinh(t) & \cosh(t) \end{pmatrix}, \begin{pmatrix} -\cosh(t) & \sinh(t) \\ \sinh(t) & -\cosh(t) \end{pmatrix}. \end{aligned}$$

With the same notation as in Theorem 1.4.1, each one of the matrices that have appeared belong to $O_1^{++}(2)$, $O_1^{--}(2)$, $O_1^{-+}(2)$ y $O_1^{+-}(2)$, respectively. We see which is the difference with the isometries of $\mathbf{E}^2$. In the latter case, and following the same scheme, it appears equations of type $x^2 + y^2 = 1$, whose solutions can write as $x = \cos \theta$ and $y = \sin \theta$. This distinguishes the equation $x^2 - y^2 = 1$, where it is necessary to separate the case that x is positive or negative.

We end this section with the study of isometries of $O_1^{++}(3)$ that leave pointwise fixed a straight-line L . This kind of isometries are called *boosts*. It will appear three types of such isometries, depending on the causal character of L .

1. L is timelike. Assume that $L = \langle E_3 \rangle$. Since $A \cdot E_3 = E_3$, we obtain that $a_{13} = a_{23} = 0$ and $a_{33} = 1$. By using the equality $G = A^t G A = G$, we have $a_{31} = a_{32} = 0$ and

$$a_{11}^2 + a_{21}^2 = 1, a_{11}a_{12} + a_{21}a_{22} = 0, a_{12}^2 + a_{22}^2 = 1.$$

Thus the matrix A writes as

$$A = \begin{pmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

2. L is spacelike. Let $L = \langle E_1 \rangle$. Then

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cosh \varphi & \sinh \varphi \\ 0 & \sinh \varphi & \cosh \varphi \end{pmatrix}.$$

3. L is lightlike. We suppose that $L = \langle E - 2 + E_3 \rangle$. Then

$$A = \begin{pmatrix} 1 & \theta & -\theta \\ -\theta & 1 - \frac{\theta^2}{2} & \frac{\theta^2}{2} \\ -\theta & -\frac{\theta^2}{2} & 1 + \frac{\theta^2}{2} \end{pmatrix}.$$

In all above cases, the isometries belong to $O_1^{++}(3)$.

We end this chapter by showing a kind of planar curves which will be used throughout this Mini-Course. Exactly these curves play the same role as the circle in Euclidean ambient. A way to define the Euclidean circle is the following. Let G be a group of rotations that leave pointwise fixed a straight-line L . Let $p_0 \notin L$. Then the set $\{A \cdot p_0; A \in G\}$ is a circle contained in a orthogonal plane to L which contains the point p_0 .

In Minkowski space $\mathbf{E}_1^3$ we have to distinguish three cases depending on the causal character of the line. Let L be a straight-line and $p_0 \notin L$. We denote by G the group of boosts associated to L . After an isometry of $\mathbf{E}_1^3$, we assume that L lies in one of the next three cases:

1. L is timelike. We consider $L = \langle E_3 \rangle$. Then

$$G = \left\{ T_\theta = \begin{pmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix}; \theta \in \mathbb{R} \right\}.$$

The set $\{T_\theta(p_0); \theta \in \mathbb{R}\}$ is the circle that lies in the plane $\{z = z_0\}$ and radius $\sqrt{x_0^2 + y_0^2}$.

2. L is spacelike. We take $L = \langle E_1 \rangle$. Then

$$G = \left\{ T_\varphi = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cosh \varphi & \sinh \varphi \\ 0 & \sinh \varphi & \cosh \varphi \end{pmatrix} \right\}.$$

The orbit of p_0 is a branch of the hyperbola $y^2 - z^2 = y_0^2 - z_0^2$ in the plane $\{x = x_0\}$.

3. L is lightlike. We assume that $L = \langle E - 2 + E_3 \rangle$. Then

$$G = \left\{ T_\theta = \begin{pmatrix} 1 & \theta & -\theta \\ -\theta & 1 - \frac{\theta^2}{2} & \frac{\theta^2}{2} \\ -\theta & -\frac{\theta^2}{2} & 1 + \frac{\theta^2}{2} \end{pmatrix}; \theta \in \mathbb{R} \right\}.$$

Let see that G leaves fixed the plane $\langle E_1, E_2 + E_3 \rangle$ since $T_\theta(E_1) = E_1 - \theta(E_2 + E_3)$. We take v the unique lightlike vector orthogonal to E_1 such that $\langle v, E_2 + E_3 \rangle = 1$. In our case, $v = E_2 - E_3$ and we consider the plane $P = \langle E_1, E_2 - E_3 \rangle$. As A is an isometry, we have that $A(E_2 - E_3)$ is a lightlike vector orthogonal to $A(E_1) = E_1 - \theta(E_2 + E_3)$ and $A(E_2 - E_3) = E_2 + E_3$. Thus it is $E_2 - E_3$. This means that $A(P) = P$. Let $p_0 \in P$. Then the set $\{T_\theta(p_0); \theta \in \mathbb{R}\}$ is a parabola included in the plane P whose axis is parallel to $E_2 - E_3$ through p_0 : see the figure 1.3. Exactly, we have

$$T_\theta(x, y, z) = (x + 2y\theta, y - x\theta - y\theta^2, -y - x\theta - y\theta^2).$$

If we set $X = x + 2y\theta$ and $Y = y - x\theta - y\theta^2$, we obtain the relation

$$Y = y + \frac{x^2}{4y} - \frac{1}{4y}X^2.$$




Figure 1.3: The orbit of a point by the action of boosts. On the left we have a hyperbola; on the right, we obtain a parabola.

We point out that the above orbits are circles, hyperbolas and parabolas *exactly* in the case that they lie in the above planes. For example, we consider the rotations with respect to the timelike line $L = \langle (0, 1, 2) \rangle$. The orbit of a point lies in a parallel plane to the the plane $P = \langle e_1 = (1, 0, 0), e_2 = (0, 2, 1) \rangle$ and it is $q + \cos(t)e_1 + \sin(t)e_2$, that is, an affine ellipse. On the other cases, we obtain affine hyperbolas and parabolas, respectively.

1.5 Exercises

1. Let U be a subspace of $\mathbb{R}^3$. Then the induced metric on U must be one of the three causal characters.
2. Let $v \in \mathbf{E}_1^3$. Then the subspace $U = \langle v \rangle$ has the same causal character than v .

3. Let $\{e_1, e_2, e_3\}$ be an orthonormal base with e_3 a timelike vector. Then any vector $v \in \mathbf{E}_1^3$ writes as

$$v = \langle v, e_1 \rangle e_1 + \langle v, e_2 \rangle e_2 - \langle v, e_3 \rangle e_3.$$

4. Let u be a spacelike vector and v a lightlike vector orthogonal to u . Show that there exists a unique lightlike vector w such that $\langle u, w \rangle = 0$ and $\langle v, w \rangle = 1$. Moreover $\{u, v, w\}$ is a base of $\mathbf{E}_1^3$ which is positive oriented and that it is future-directed.
5. In the set $\mathcal{T}$ of timelike vectors we define the following relations: $u \sim v$ if $\langle u, v \rangle < 0$. Without using the concept of timelike cone, show that $\sim$ is a equivalence relation with only two classes, which agrees with the sets of future-directed vectors and past-directed vectors. If we put $u \sim v$ if $\langle u, v \rangle > 0$, then $\sim$ is not an equivalence relation.
6. Let B and B' be two orthonormal bases and let $A = (a_{ij})$ the matrix of change of bases. Show that B and B' belong to the same timelike orientation if $a_{33} > 0$.
7. Let $u, v \in \mathbf{E}_1^3$ be two independent linear vectors. Show that if u and v are spacelike (resp. timelike, lightlike) vectors, then $u \times v$ is a timelike (resp. spacelike, spacelike) vector. If one is timelike and the other one is lightlike, then, $u \times v$ is spacelike. Finally, if u is spacelike and v is lightlike, then $u \times v$ can be spacelike or lightlike.
8. Let $B = \{e_1, e_2, e_3\}$ be an orthonormal base of $\mathbf{E}_1^3$. Show that if u and v are two vectors with coordinates (u_1, u_2, u_3) and (v_1, v_2, v_3) respectively, then the coordinates of $u \times v$ agrees with the expression (1.3).
9. Let $u, v \in \mathcal{T}$ be two vectors in the same timelike cone. Then

$$|u \times v|^2 = |u|^2 |v|^2 \sinh(\varphi)^2, \quad \varphi = \text{angle}(u, v).$$

10. Let $u \neq 0$. Then

$$|u| = |u|_\epsilon \sqrt{|\cos(2\theta)|} \leq |u|_\epsilon,$$

where θ is the angle of u with any horizontal plane. Equality holds if and only if u is horizontal or vertical.

11. If u and v are independent linear, then

$$\langle u \times v, u \times v \rangle = \langle u \times_\epsilon v, u \times_\epsilon v \rangle_\epsilon \cos(2\theta).$$

12. $|u \times v| \leq |u \times_\epsilon v|_\epsilon$. Equality holds if and only if $\{u, v\}$ determine a horizontal or vertical plane.
13. $u \times v$ y $u \times_\epsilon v$ lies in the same halfspace (resp. opposite halfspace) determined by the plane $\langle u, v \rangle$ when $u \times v$ is a timelike (resp. spacelike) vector.
14. Let A be a isometry of $\mathbf{E}_1^3$ and $u, v \in \mathbf{E}_1^3$. Then $A(u \times v) = \det(A)(Au) \times (Av)$.

15. If u and v are vectors of $\mathbf{E}_1^3$, we denote by $[u, v]$ the segment joining both vectors, that is, the set $\{tu + (1-t)v; t \in [0, 1]\}$. Show that if u and v are spacelike, then all vectors of $[u, v]$ are spacelike. This does not occur if one vector is spacelike and the other one is timelike.

16. Find the isometries that leave pointwise fixed the y -axis.

17. Show that the causal character of a vector or subspace does not change if we map it by a isometry of $O_1^{++}(3)$. This does not occur if the isometry is of other kind.
18. Show that any element of $O_1^{++}(3)$, except the identity, pointwise fixed leaves exactly one line. This shows that any element of $O^{++}(3)$ is a boost.
19. Let $\{u_i\}, \{v_i\}$ be two bases of $\mathbf{E}_1^3$ that satisfy that u_1 and v_1 are two unit spacelike vectors, u_i, v_i are lightlike vectors, $j \in \{2, 3\}$ and

$$\langle u_1, u_j \rangle = \langle v_1, v_j \rangle = 0, \quad j \in \{2, 3\}$$

$$\langle u_2, u_3 \rangle = \langle v_2, v_3 \rangle = 1.$$

Assume that both bases are positive oriented and future-directed. Show that there exists $A \in O_1^{++}(3)$ that carries a base in the other one.

20. Show that $A \in O_1^+(3)$ if and only $a_{33} > 0$.

Chapter 2

Curves in Minkowski space

In this chapter we develop the theory of the Frenet trihedron for curves in $\mathbf{E}_1^3$. In this sense, we treat similar questions to what happen in Euclidean ambient. For example, we will find all planar curves with constant curvature. Also, we will study helices and Bertrand curves.

The first point to remark is that the notion of curve is not metric: a (smooth) curve is a differentiable map $\alpha : I \rightarrow \mathbb{R}^3$ where I is an open of $\mathbb{R}$. It is the fact to consider on I the induced metric by α what converts into a differential geometric concept. The development to do is the same than for curves in Euclidean curves in $\mathbf{E}^3$. However, the different causal character that can have a straight-line in $\mathbf{E}_1^3$ will do that its study is more difficult because we have to consider each causal character.

In this chapter $\alpha : I \subset \mathbb{R} \rightarrow \mathbf{E}_1^3$ denotes a differentiable map, where I is an open interval, with $0 \in I$.

2.1 Parametrized curves

Definition 2.1.1. *Let α be a curve in $\mathbf{E}_1^3$. We say that α is spacelike (resp. timelike, lightlike) at t if $\alpha'(t)$ is a spacelike (resp. timelike, lightlike) vector. The curve α is called spacelike (resp. timelike, lightlike) if it is for any $t \in I$.*

Before to show examples of curves, we remark that in general any curve in $\mathbf{E}_1^3$ is not of one type of the above ones. Of course, for each $t \in I$, $\alpha'(t)$ will be spacelike, timelike or lightlike, but this property does not maintain in all interval I . For example, if we consider the curve $\alpha(t) = (\cosh(t), t^2, \sinh(t))$, then $\langle \alpha'(t), \alpha'(t) \rangle = 4t^2 - 1$. Thus, the curve is spacelike in the interval $(-\infty, -1/2) \cup (1/2, \infty)$, timelike in $(-1/2, 1/2)$ and lightlike in $\{-1/2, 1/2\}$.

We point out that the spacelike (or timelike) condition is an open property, that is, if α is spacelike (or timelike) at $t_0 \in I$, there exists an interval $(t_0 - \delta, t_0 + \delta)$ where α has the same spacelike (or timelike) character: if at $t_0 \in I$ we have $\langle \alpha'(t_0), \alpha'(t_0) \rangle > 0$ (< 0), the continuity assures the existence of an interval around t_0 where $\langle \alpha'(t_0), \alpha'(t) \rangle > 0$ (< 0).

A natural way to justify the definition of the causal character is the following. Let $\alpha : I \rightarrow \mathbf{E}_1^3$ a

differentiable curve. For each $t \in I$, we consider the differential map $(d\alpha)_t : T_t I \equiv \mathbb{R} \rightarrow T_{\alpha(t)} \mathbf{E}_1^3 \equiv \mathbb{R}^3$, that is, the map given by

$$(d\alpha)_t(s) = \frac{d}{du}|_{u=0} \alpha(t + su) = s \cdot \alpha'(t),$$

or,

$$(d\alpha)_t = \alpha'(t).$$

Let us observe that $(d\alpha)_t(\frac{\partial}{\partial t}) = \alpha'(t)$. We consider in $\mathbb{R} = T_t I$ the induced metric from α , that is, $\alpha^*\langle, \rangle$ which is defined by

$$\alpha^*\langle, \rangle_t(m, n) = \langle (d\alpha)_t(m), (d\alpha)_t(n) \rangle = mn \langle \alpha'(t), \alpha'(t) \rangle.$$

If we take the usual base at $T_t I$, that is, $\frac{\partial}{\partial t}$, we have

$$\alpha^*\langle, \rangle_t(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}) = \langle \alpha'(t), \alpha'(t) \rangle.$$

Then $(T_t I, \alpha^*\langle, \rangle)$ is a metric space of dimension one. We now rediscover the causal character of α at t : it agrees with the metric space $(T_t I, \alpha^*\langle, \rangle)$, that is, if this space is positive definite, or $\alpha'(t) = 0$, we say that α is spacelike at t ; if the space is negative definite, then α is timelike at t ; and if the space is degenerate, we say that α is lightlike at t .

A curve α is called *regular* at $t_0 \in I$ if $\alpha'(t_0) \neq 0$. We say that α is regular if it is for any $t \in I$.

We show some examples of planar curves in $\mathbf{E}_1^3$, that is, curves included in an affine plane of $\mathbb{R}^3$. Let $p, v \in \mathbb{R}^3$ and $r > 0$.

1. The straight-line $\alpha(t) = p + tv$ has the same causal character than v .
2. The circle $\alpha(t) = p + r(\cos t, \sin t, 0)$ is a spacelike curve included in a spacelike plane.
3. The hyperbola $\alpha(t) = p + r(0, \sinh t, \cosh t)$ is a spacelike curve in a timelike plane.
4. The hyperbola $\alpha(t) = p + r(0, \cosh t, \sinh t)$ is timelike in a lightlike plane.
5. The parabola $\alpha(t) = (t, t^2, t^2)$ is a spacelike curve in a lightlike plane.

Let now see examples of non-planar curves.

1. The helix $\alpha(t) = (\cos t, \sin t, at)$, $a \neq 0$. This curve is an Euclidean helix.
2. $\alpha(t) = (at, \sinh t, \cosh t)$, $a \neq 0$.
3. $\alpha(t) = (at, \cosh t, \sinh t)$, $a \neq 0$.

The causal character of a curve in Minkowski space imposes conditions on the regularity and topology of the curves. First, we have

Proposition 2.1.2. *Any timelike or lightlike curve is regular.*

Proof: Assume that the curve is timelike, and we write $\alpha(t) = (x(t), y(t), z(t))$, where the function x , y and z are differentiable functions on t . Then $\langle \alpha'(t), \alpha'(t) \rangle = x'(t)^2 + y'(t)^2 - z'(t)^2 < 0$, in particular, $z'(t) \neq 0$, that is, α is a regular curve.

If the curve is lightlike, we have $z'(t) \neq 0$ again since, on the contrary, $x'(t) = y'(t) = 0$ and $\alpha'(t) = 0$. But this means that α is spacelike at t . $\square$

As a consequence of the proof, around t_0 any timelike or lightlike curve can locally written as $\alpha(t) = (f(t), g(t), t)$, $t \in (t_0 - \delta, t_0 + \delta)$ for some smooth functions f and g . For a spacelike curve, we have the following result: given $t_0 \in I$, there exists $\delta > 0$ such that $\alpha(t) = (t, f(t), g(t))$ or $\alpha(t) = (f(t), t, g(t))$.

Other consequence about the causal character is the following

Theorem 2.1.3. *Let α be a closed curve in $\mathbf{E}_1^3$ included in an affine plane P .*

1. *If α is spacelike, then P is a spacelike plane.*
2. *The curve is not timelike or lightlike.*

Proof:

1. Without loss of generality, we assume that the plane P is a vector plane. If P is timelike, we assume that $P = \langle E_2, E_3 \rangle$. Then $\alpha(t) = (0, y(t), z(t))$. The map given by $y : \mathbb{R} \rightarrow \mathbb{R}$ attains a maximum at some point t_0 . Then $y'(t_0) = 0$, and $\alpha'(t_0) = (0, 0, z'(t_0))$ is a timelike vector.
If P is a lightlike plane, we take $P = \langle E_1, E_2 + E_3 \rangle$. Then $\alpha(t) = (x(t), y(t), y(t))$. Let t_0 be the maximum of the function $x(t)$. Then $\alpha'(t_0) = (0, y'(t_0), y'(t_0))$ is a lightlike vector: contradiction. As a consequence, P is a spacelike plane.
2. Let α be a timelike curve. Then the plane has to be timelike since in spacelike or lightlike planes there are not timelike vectors. If $P = \langle E_2, E_3 \rangle$, the point t_0 where the function $z(t)$ attains the maximum satisfies $\alpha'(t_0) = (0, y'(t_0), 0)$: this vector is spacelike. This is a contradiction. An analogous reasoning can do for lightlike curves.

$\square$

With the same arguments, we have

Corollary 2.1.4. *There are not closed curves in $\mathbf{E}_1^3$ that are timelike or lightlike.*

Let us see that there exist (non closed) spacelike curves in non-spacelike planes. As an example, let $\alpha(s) = (0, \sinh(s), \cosh(s))$: this is a spacelike curve in the timelike plane $\langle E_2, E_3 \rangle$. Similarly, the curve $\alpha(s) = (s, s^2, s^2)$ is spacelike and included in the lightlike plane $\langle E_1, E_2 + E_3 \rangle$.

From now, we will assume that all curves are regular.

Lemma 2.1.5. *Let α be a spacelike or timelike curve. Then there exists a change of parameters such that $|\alpha'(s)| = 1$. Exactly, given $t_0 \in I$, there is $\delta, \epsilon > 0$ and a diffeomorphism $\phi : (-\epsilon, \epsilon) \rightarrow (t_0 - \delta, t_0 + \delta)$ such that the curve $\beta : (-\epsilon, \epsilon) \rightarrow \mathbf{E}_1^3$ given by $\beta = \alpha \circ \phi$ satisfies the property $|\beta'(s)| = 1$. In such case, we say that the curve is arc-length parameterized.*

Proof: We do the proof for timelike curves. We define the map $S : I \rightarrow \mathbb{R}$ by

$$S(t) = - \int_{t_0}^t \langle \alpha'(u), \alpha'(u) \rangle du.$$

Since $S'(t_0) > 0$, the function S is a local diffeomorphism around $t = t_0$. Because $S(t_0) = 0$, there exist $\delta, \epsilon > 0$ such that $S : (t_0 - \delta, t_0 + \delta) \rightarrow (-\epsilon, \epsilon)$ is a diffeomorphism. The map that we are looking for is $\phi = S^{-1}$. $\square$

For lightlike curves, the vector $\alpha'(t)$ is lightlike and there would be not sense re-parametrize by the arc-length. However and as $\langle \alpha'(t), \alpha'(t) \rangle = 0$, by differentiation, we obtain $\langle \alpha''(t), \alpha'(t) \rangle = 0$. Let us suppose that $\alpha''(t) \neq 0$. Then $\alpha''(t)$ is a spacelike vector and we can parametrized α to get $|\alpha''(t)| = 1$. The following result asserts that this can be done.

Lemma 2.1.6. *Let α be a lightlike curve in $\mathbf{E}_1^3$. There exists a re-parametrization of α given by $\beta(s) = \alpha(\phi(s))$ in such way that $|\beta''(s)| = 1$. We say that α is length-arc pseudo-parametrized.*

Proof: We write $\beta(s) = \alpha(\phi(s))$, where the function ϕ is unknown. By twice differentiation, we have

$$\beta''(s) = \phi''(s)\alpha'(t) + \phi'(s)^2\alpha''(t).$$

Then $\langle \beta'(s), \beta'(s) \rangle = \phi'(s)^4 |\alpha''(t)|^2$. Thus, we define ϕ as the solution of the differential equation

$$\phi'(s) = \frac{1}{\sqrt{|\alpha''(\phi(s))|}}, \quad \phi(0) = t_0.$$

$\square$

2.2 Curvature and torsion

This section is the key of this chapter. Given a regular curve, we want to assign for any point of the curve an orthonormal base that describes the geometry of the curve. This will be given by the Frenet trihedron. The variation of this base along the curve will give us the information how the curve is deformed into the ambient space.

The simplest case of curves are straight-lines. If $p \in \mathbf{E}_1^3$ and $v \neq 0$, the straight-line through the point p in the direction v is parametrized by $\alpha(t) = p + tv$. Then $\alpha''(t) = 0$. As the acceleration¹ has modulus 0, we say that the *curvature* of a straight-line is 0.

Conversely, if α is a regular curve that satisfies $\alpha''(s) = 0$ for any s , by integration twice, it yields $\alpha'(s) = v$, for some vector $v \neq 0$ and $\alpha(s) = sv + p$. This means that α parametrizes a straight-line through the point p along the direction given by v . Let us see that given a straight-line (as a set of $\mathbb{R}^3$), we can parametrize it by other parametrization. For example, $\alpha(s) = s^3 E_1$ is a parametrization of the straight-line $\langle E_1 \rangle$ and does not satisfy $\alpha'' = 0$.

We consider a regular curve α parametrized by the length-arc or the by the pseudo-length-arc. We call $\mathbf{T}(s) = \alpha'(s)$ the *tangent vector* at s . In particular, $\langle \mathbf{T}(s), \mathbf{T}'(s) \rangle = 0$. We will assume that $\mathbf{T}'(s) \neq 0$ and that $\mathbf{T}'(s)$ is not proportional to $\mathbf{T}(s)$ for each s . This avoids that the curve is a straight-line.

¹We avoid to say that the acceleration is null

2.2.1 The timelike case

We suppose that α is a timelike curve. Then $\mathbf{T}'(s) \neq 0$ is a spacelike vector independent with $\mathbf{T}(s)$. We define the *curvature* of α at s as $\kappa(s) = |\mathbf{T}'(s)|$. The *normal vector* $\mathbf{N}(s)$ is defined by

$$\mathbf{N}(s) = \frac{\mathbf{T}'(s)}{\kappa(s)} = \frac{\alpha''(s)}{|\alpha''(s)|}.$$

Moreover $\kappa(s) = \langle \mathbf{T}'(s), \mathbf{N}(s) \rangle$. We call the *binormal vector* $\mathbf{B}(s)$ as

$$\mathbf{B}(s) = \mathbf{T}(s) \times \mathbf{N}(s).$$

The vector $\mathbf{B}(s)$ is unitary and spacelike. For each s , $\{\mathbf{T}, \mathbf{N}, \mathbf{B}\}$ is an orthonormal base of $\mathbf{E}_1^3$ which is called the *Frenet trihedron* of α . We define the torsion of α at s as

$$\tau(s) = \langle \mathbf{N}'(s), \mathbf{B}(s) \rangle.$$

By differentiation each one of the vector functions of the Frenet trihedron and putting in relation with the same Frenet base, we obtain the Frenet equations, namely,

$$\begin{pmatrix} \mathbf{T}' \\ \mathbf{N}' \\ \mathbf{B}' \end{pmatrix} = \begin{pmatrix} 0 & \kappa & 0 \\ \kappa & 0 & \tau \\ 0 & -\tau & 0 \end{pmatrix} \begin{pmatrix} \mathbf{T} \\ \mathbf{N} \\ \mathbf{B} \end{pmatrix}.$$

2.2.2 The spacelike case

Let α be a spacelike curve. There are three possibilities depending on the causal character of $\mathbf{T}'(s)$.

1. The vector $\mathbf{T}'(s)$ is spacelike. Again, we write $\kappa(s) = |\mathbf{T}'(s)|$, $\mathbf{N}(s) = \mathbf{T}'(s)/\kappa(s)$ and $\mathbf{B}(s) = \mathbf{T}(s) \times \mathbf{N}(s)$. The vectors $\mathbf{N}$ and $\mathbf{B}$ are called the normal vector and the binormal vector respectively. The curvature of α is defined by κ . The Frenet equations are

$$\begin{pmatrix} \mathbf{T}' \\ \mathbf{N}' \\ \mathbf{B}' \end{pmatrix} = \begin{pmatrix} 0 & \kappa & 0 \\ -\kappa & 0 & \tau \\ 0 & \tau & 0 \end{pmatrix} \begin{pmatrix} \mathbf{T} \\ \mathbf{N} \\ \mathbf{B} \end{pmatrix}.$$

The torsion of α is defined by $\tau = -\langle \mathbf{N}', \mathbf{B} \rangle$.

2. The vector $\mathbf{T}'(s)$ is timelike. The normal vector is $\mathbf{N} = \mathbf{T}'/\kappa$, where $\kappa(s) = \sqrt{-\langle \mathbf{T}'(s), \mathbf{T}'(s) \rangle}$ is the curvature of α . The binormal vector is $\mathbf{B} = \mathbf{T} \times \mathbf{N}$, which is a spacelike vector. Now, the Frenet equations are

$$\begin{pmatrix} \mathbf{T}' \\ \mathbf{N}' \\ \mathbf{B}' \end{pmatrix} = \begin{pmatrix} 0 & \kappa & 0 \\ \kappa & 0 & \tau \\ 0 & \tau & 0 \end{pmatrix} \begin{pmatrix} \mathbf{T} \\ \mathbf{N} \\ \mathbf{B} \end{pmatrix}.$$

The torsion of α is $\tau = \langle \mathbf{N}', \mathbf{B} \rangle$.

3. The vector $\mathbf{T}'(s)$ is lightlike for any s (recall that $\mathbf{T}'(s) \neq 0$ and it is not proportional to $\mathbf{T}(s)$). We define the normal vector as $\mathbf{N}(s) = \mathbf{T}'(s)$, which is independent linear with $\mathbf{T}(s)$. Let $\mathbf{B}$ be the unique lightlike vector such that $\langle \mathbf{N}, \mathbf{B} \rangle = 1$ and orthogonal to $\mathbf{T}$. The vector $\mathbf{B}(s)$ is called the binormal vector of α at s . The Frenet equations are

$$\begin{pmatrix} \mathbf{T}' \\ \mathbf{N}' \\ \mathbf{B}' \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & \tau & 0 \\ -1 & 0 & -\tau \end{pmatrix} \begin{pmatrix} \mathbf{T} \\ \mathbf{N} \\ \mathbf{B} \end{pmatrix}.$$

The function τ is called the torsion of α . There is not a definition of the curvature of α .

2.2.3 The lightlike case

Let α be a lightlike curve parametrized by the pseudo-length-arc, that is, $\alpha''(s)$ is a unitary spacelike vector. We consider the tangent vector $\mathbf{T} = \alpha'$ and we define the normal vector as $\mathbf{N}(s) = \mathbf{T}'(s)$ and the binormal vector the unique lightlike vector orthogonal to $\mathbf{N}(s)$ such that $\langle \mathbf{N}(s), \mathbf{B}(s) \rangle = 1$. Then the Frenet equations are:

$$\begin{pmatrix} \mathbf{T}' \\ \mathbf{N}' \\ \mathbf{B}' \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ \tau & 0 & -1 \\ 0 & -\tau & 0 \end{pmatrix} \begin{pmatrix} \mathbf{T} \\ \mathbf{N} \\ \mathbf{B} \end{pmatrix}.$$

Again, the function τ is the torsion of α . We point out that as in the case that α is spacelike with $\mathbf{T}'$ lightlike, we do not define the curvature of the curve.

We end this section with some final remarks on the general theory of curves. One can prove that the notions of curvature and torsion are isometric invariant, that is, if $M : \mathbf{E}_1^3 \rightarrow \mathbf{E}_1^3$ is a rigid motion of $\mathbf{E}_1^3$, and we consider the curve $\beta(s) = M \circ \alpha(s)$, then $\kappa_\beta(s) = \kappa_\alpha(s)$ and $\tau_\beta(s) = \pm \tau_\alpha(s)$.

Similarly as in Euclidean ambient, one can find formulae for the curvature and torsion function in the case that the curve is not parametrized by the length-arc. First, we need to define the curvature and torsion for this kind of curves. Let $\beta = \alpha \circ \phi$ be any parametrization by the length-arc. We define $\kappa_\alpha(t) = \kappa_\beta \circ \phi^{-1}$ and similar for the torsion. One can show that this definition does not depend on the re-parametrization. For example, for timelike curves, we have:

$$\kappa_\alpha(t) = \frac{|\alpha'(t) \times \alpha''(t)|}{|\alpha'(t)|^3} = \frac{|\alpha'(t) \times \alpha''(t)|}{(-\langle \alpha'(t), \alpha'(t) \rangle)^{3/2}}$$

$$\tau(t) = \frac{\det(\alpha'(t), \alpha''(t), \alpha'''(t))}{|\alpha'(t) \times \alpha''(t)|^2}.$$

In the case of planar curves, we have the following result for timelike curves and which is analogous to one for Euclidean planar curves.

Theorem 2.2.1. *Let $\kappa : I \rightarrow \mathbb{R}$ be a smooth function and let P a timelike plane. There exists a unique spacelike curve in P with curvature κ . The same result holds to assure the existence of a timelike curve in P with κ as curvature function.*

Proof: Without loss of generality, we assume that $P = \langle E_1, E_3 \rangle$ and let $p_0 = (0, 0, 0)$.

1. We are going to find the spacelike curve α such that $\alpha(0) = (0, 0, 0)$, $\alpha'(0) = E_1$ and $\kappa_\alpha(s) = \kappa(s)$. Let $\theta : I \rightarrow \mathbb{R}$ be the function

$$\theta(s) = \int_0^s \kappa(t) dt,$$

and

$$x(s) = \int_0^s \cosh(\theta(t)) dt, \quad z(s) = \int_0^s \sinh(\theta(t)) dt.$$

Then $\alpha(s) = (x(s), z(s))$ is the curve that we are looking for. First, we have $\alpha(0) = (0, 0)$. As

$$\alpha'(s) = (\cosh(\theta(s)), \sinh(\theta(s))) \quad \alpha''(s) = \kappa(s)(\sinh(\theta(s)), \cosh(\theta(s))),$$

then α is a parametrized by the length-arc spacelike curve with $\alpha'(0) = (\cosh(0), \sinh(0)) = (1, 0)$. Therefore the curvature of α is $|\alpha''(s)|$, that is, $\kappa(s)$.

2. If we look for a timelike curve with starting velocity E_3 , it suffices to take $\theta(s) = \int_0^s \kappa(t) dt$ and

$$x(s) = \int_0^s \sinh(\theta(t)) dt, \quad z(s) = \int_0^s \cosh(\theta(t)) dt.$$

□

2.3 Planar curves with constant curvature

We consider a planar curve, that is, a curve of $\mathbf{E}_1^3$ included in a affine plane. After a rigid motion, we can assume that this plane is a vector plane. In this section we are going to study *planar* curves with *constant curvature*. Before, we translate to the Lorentzian ambient a result that holds for curves in a Euclidean plane. Recall that if α is a planar curve in $\mathbf{E}^2$ parametrized by the length-arc and v is a fixed unitary direction, we call $\theta(s)$ the angle that makes $\mathbf{T}(s)$ and v , that is, $\cos(\theta(s)) = \langle \mathbf{T}(s), v \rangle$. One can prove that the curvature of α is $|\theta'(s)|$.

In the Lorentzian ambient, we can not extend this result since it is necessary to talk of angle². We only do it for timelike curves.

Theorem 2.3.1. *Let α be a timelike curve parametrized by the length-arc and included in a timelike plane. Let v be a unit fixed vector of the plane pointing to the future. Let $\phi(s)$ be the hyperbolic angle between $\mathbf{T}(s)$ and v . Then $\kappa(s) = |\phi'(s)|$.*

Proof: Without loss of generality, we take $P = \langle E_1, E_2 \rangle$ and $v = (0, v_2, v_3)$ with $v_3 > 0$. Recall that

$$-\cosh \phi(s) = \langle \mathbf{T}(s), v \rangle.$$

By differentiation, we obtain

$$-\phi'(s) \sinh \phi(s) = \kappa(s) \langle \mathbf{N}(s), v \rangle.$$

²See the corresponding exercise at the end of this chapter

As $\{\mathbf{N}, \mathbf{T}\}$ is a base of P , then

$$v = \langle v, \mathbf{N}(s) \rangle \mathbf{N}(s) - \langle v, \mathbf{T}(s) \rangle \mathbf{T}(s).$$

Then

$$-1 = \langle v, \mathbf{N}(s) \rangle^2 - \langle v, \mathbf{T}(s) \rangle^2 = \langle v, \mathbf{N}(s) \rangle^2 - \cosh(\phi(s))^2,$$

that is, $\langle v, \mathbf{N}(s) \rangle = \pm \sinh(\phi(s))$. Therefore $|\phi'(s)| = \kappa(s)$. □

Once showed this result, we are going to study planar curves with constant curvature. We will assume that this constant is not 0 (if it is, we know that α is a straight-line). The study of these curves has to distinguish the causal character of the curves. We will re-discover the curves that appeared in Chapter 1 as orbits of a point by the motion of a group of boosts (isometries that pointwise fixed leave a straight-line).

1. **(The timelike case)** The plane containing the curve must be timelike. Thus we assume that it is $P = \langle E_2, E_3 \rangle$ and so, $\alpha(s) = y(s)E_2 + z(s)E_3$. Because the curve is parametrized by the length-arc, $y'(s)^2 - z'(s)^2 = -1$. Then

$$z'(s) = \cosh \phi(s), \quad y'(s) = \sinh \phi(s).$$

We compute the curvature of α :

$$\kappa(s) = |\alpha''(s)| = |\phi'(s)| := a.$$

Then $\phi(s) = as + b$. It yields

$$y' = \sinh(As + b), \quad z'(s) = \cosh(as + b).$$

Then

$$\alpha(s) = \frac{1}{a}(\cosh(as + b)E_2 + \sinh(as + b)E_3).$$

This curve is a Euclidean hyperbola in the plane P .

2. **(The spacelike case)** We are going to find the planar spacelike curves with constant curvature.
 - (a) Let us assume that $P = \langle E_1, E_2 \rangle$. The induced metric on P by the Lorentzian metric agrees with the Euclidean metric. Thus, in this situation³, the notions of curvature agree in both settings. In particular, α is an Euclidean circle.
 - (b) If the plane is timelike, we take the plane given by $\{x = 0\}$. We write α as $\alpha(s) = (x(s), 0, z(s))$, with $x'(s)^2 - z'(s)^2 = 1$. Thus, there is a function $\theta \in \mathbb{R}$ such that $x'(s) = \cosh(\theta(s))$ y $z'(s) = \sinh(\theta(s))$. An analogous reasoning as above yields

$$\alpha(s) = (x(s), z(s)) = \left(\frac{1}{a} \sinh(as + b), \frac{1}{a} \cosh(as + b)\right) = \frac{1}{a}(\sinh(as + b), \cosh(as + b)).$$

This curve is an Euclidean hyperbola.

³Let see the corresponding exercise that shows that even the plane is spacelike, the Euclidean and Lorentzian curvature do not agree

- (c) We suppose that the plane is lightlike. In this case there is not notion of curvature of a curve. After a rigid motion, we assume that the plane containing the curve is $\{y - z = 0\}$. Now the curve writes as $\alpha(s) = (x(s), y(s), y(s))$. As α is a spacelike curve which it is parametrized by the length-arc, then $x'(s)^2 = 1$. Let us assume $x(s) = s$. Then $\mathbf{T}(s) = (1, y'(s), y'(s))$. We know that $\mathbf{T}'(s)$ is a lightlike vector. As in the plane there is a unique lightlike direction, then the vector $\mathbf{T}'(s)$ is proportional to a fixed vector, for example, to $v = (0, 1, 1)$. We say that α has constant curvature if $\tau = 0$. As a consequence, $\mathbf{T}'(s) = v$ (up constants). By integration, we obtain $\alpha(s) = p_0 + sw + \frac{s^2}{2}(E_2 + E_3)$, where w is a unit spacelike vector. As $\langle w, w \rangle = 1$, the vector w writes as $w = E_1 + c(E_2 + E_3)$ with $c \in \mathbb{R}$. Then

$$\alpha(s) = p_0 + sE_1 + \left(cs + \frac{s^2}{2}\right)(E_2 + E_3),$$

that is, and up re-parametrization, it is a parabola in the plane P with axis parallel to the lightlike direction.

3. **(The lightlike case)** Let α be a lightlike curve included in a plane P . The plane P can be lightlike or timelike. We separate both situations:

- (a) The plane is lightlike. Assume that $P = \langle E_1, E_2 + E_3 \rangle$. As $\alpha'(s)$ is lightlike and there is only one lightlike directions, then $\alpha'(s) = f(s)(E_2 + E_3)$ for some differentiable function f . In such case, α is a straight-line. Recall that this case was not considered in the Frenet trihedron.
- (b) The plane is timelike. Let $P = \langle E_2, E_3 \rangle$. As there are only two directions and both are independent linear, namely, $E_2 + E_3$ and $E_2 - E_3$, and $\alpha'(s) = f(s)(E_2 \pm E_3)$, an argument of connection together the fact that the vector $v = 0$ is not lightlike, shows that either $\alpha'(s) = f(s)(E_2 + E_3)$ in I or $\alpha'(s) = f(s)(E_2 - E_3)$ in I . Anyway, the curve is a straight-line.

If we do realize a similar argument to the case that α is spacelike and $\mathbf{N}$ is lightlike, one could say that the curve has constant curvature if $\tau = 0$ in I . Then $\mathbf{N}'(s) = 0$. By integration, we get $\alpha'(s) = sv + w$, where $\mathbf{N}(s) = v$ is a spacelike vector. But as $\langle \mathbf{T}, \mathbf{N} \rangle = 0$, then $s + \langle v, w \rangle = 0$ for each s : contradiction. The only possibility is that $v = 0$, that is, α is a straight-line.

Theorem 2.3.2. *After a rigid motion of $\mathbf{E}_1^3$, the only planar curves with (non-zero) constant curvature are Euclidean circles, hyperbolas and parabolas.*

2.4 Helices and Bertrand curves in $\mathbf{E}_1^3$

In Euclidean space, a helix is a curve whose tangent straight-lines make a constant angle with a fixed direction. This direction is called the axis of the helix. A result due to Lancret shows that a curve is a helix if and only if τ/κ is a constant function. For example, planar curves are helices. A helix with constant curvature and torsion is called a *cylindrical helix*.

We extend this notion to the Lorentzian ambient. Although we can not speak about the angle between vectors, we can consider the function $\langle \mathbf{T}, v \rangle$ for a fixed direction v and to say that this function is constant. In this sense, we have:

Definition 2.4.1. A helix in $\mathbf{E}_1^3$ is a regular curve such that $\langle \mathbf{T}(s), v \rangle$ is a constant function for some fixed vector $v \neq 0$. Any line parallel this direction v is called the axis of the helix.

We assume that the curve is not planar.

Theorem 2.4.2. If a (timelike or spacelike) curve α in $\mathbf{E}_1^3$ is a helix, then τ/κ is a constant function.

Proof: We distinguish the cases that α is spacelike and timelike.

- Let α be a spacelike curve. There are three possibilities.
 - $\mathbf{T}'$ is a spacelike vector. By differentiation, $\langle \mathbf{T}, v \rangle = ct$ we obtain $\kappa \langle \mathbf{N}, v \rangle = 0$. Then, $\langle \mathbf{N}, v \rangle = 0$. There exist functions a, b such that $v = a\mathbf{T} + b\mathbf{B}$. By differentiation with respect to s and using the Frenet equations, we get

$$a' = b' = 0, \quad a\kappa + b\tau = 0,$$

obtaining the desired result.

- If $\mathbf{T}'$ is a timelike vector, the reasoning is analogous.
- Let $\mathbf{T}'$ be a lightlike vector. The same reasoning says us that

$$0 = (a' - b)\mathbf{T} + a\mathbf{N} + (b' - \tau b)\mathbf{B},$$

which implies that $a = b = 0$, that is, $v = 0$: contradiction.

- The case that α is timelike is similar to the the case that α and $\mathbf{N}$ are both spacelike.

□

In order to say the converse of this theorem, we need the notions of curvature and torsion. For this reason, we have:

Theorem 2.4.3. Let α be a timelike curve or a spacelike curve with non-lightlike normal vector. If τ/κ is constant, then α is a helix.

Proof: Assume that $\tau = c\kappa$, $c \in \mathbb{R}$. We are going only to study the case that α is timelike. We define the vector function given by

$$v(s) = c\mathbf{T}(s) + \mathbf{B}(s).$$

By using the Frenet equations, we show that $v' = 0$, that is, v is a constant function. Moreover, $\langle \mathbf{T}, v \rangle = c = ct$, which proves that α is a helix. □

One can think what does happen in the cases that α is spacelike with normal vector lightlike, or in the case that α is a lightlike curve. We consider the first case. We point out that any planar curve is a helix: we take $P = \langle E_1, E_2 + E_3 \rangle$; since $\alpha'(s)$ belongs to this plane and $E_2 + E_3 \in P^\perp$, then $\langle \mathbf{T}(s), E_2 + E_3 \rangle = 0$.

On the other had, if α satisfies $\langle \mathbf{T}(s), v \rangle = a$, by using the Frenet equations, the vector v writes as $v = a\mathbf{T} + b(s)\mathbf{N}(s)$. By differentiation, we obtain $b' + b\tau + a = 0$. Let us see that any spacelike

curve with normal vector of lightlike causal character is a helix. Let a be any constant and $b = b(s)$ a solution of the differential equation $b'(s) + \tau(s)b(s) + a = 0$. We define $v(s) = a\mathbf{T}(s) + b(s)\mathbf{N}(s)$. This function is constant since $v'(s) = (b' + b\tau + a)\mathbf{N} = 0$. Moreover $\langle \mathbf{T}, v \rangle = a$, and so, α is a helix. In this situation, there exists an infinity set of vectors v .

In Euclidean ambient, a *Bertrand curve* is a regular curve such that there exists other curve β whose straight-lines in the corresponding points are parallel. The curve β is called the *mate curve* of α . Therefore, the planar curves are Bertrand: it suffices to take parallel curves to the original curve. If $\tau \neq 0$, a characterization of a Bertrand curve is the following: the curve α is a Bertrand curve if and only if $A\kappa + B\tau = 1$ for some constant A and B (if $A\kappa + B\tau = 0$, we have an helix). As example of Bertrand curves are the cylindrical helices, that is, curves where κ and τ are constant functions.

In $\mathbf{E}_1^3$ we have the same definition of a Bertrand curve as in Euclidean setting. We focus our attention for timelike curves.

Theorem 2.4.4. *Let α be a timelike curve in $\mathbf{E}_1^3$. Then α is a Bertrand curve if and only if there are two constants A and B such that*

$$A\kappa + B\tau = 1.$$

Proof: If α is a Bertrand curve, let β be the mate curve of α . Then β can parametrize as $\beta(s) = \alpha(s) + \lambda(s)\mathbf{N}(s)$. By using the Frenet equations, we have

$$\beta'(s) = (1 + \lambda\kappa)\mathbf{T} + \lambda'\mathbf{N} + \lambda\tau\mathbf{B}.$$

Thus $\lambda' = 0$, that is, λ is a constant function.

Assume that β is a lightlike curve. Then $1 + \lambda\kappa = \pm\lambda\tau$ and we have the result. Let now the case that β is a non-degenerate curve, for example, a spacelike curve. We find the normal direction of β . For this, we parametrize β by the length-arc: $\gamma(s) = \beta(\phi(s))$. Then the normal direction of β is given by $\gamma''(s)$. We have $\gamma'(s) = \phi'(s)\beta'(t)$ and

$$\gamma''(s) = \phi''\beta' + \phi'^2\beta''.$$

Now

$$\begin{aligned} \phi'^2 &= \frac{1}{\langle \beta', \beta' \rangle} \\ \phi'' &= -\frac{\langle \beta'', \beta' \rangle}{\langle \beta', \beta' \rangle^2}. \end{aligned}$$

We then have an expression of γ'' in terms of the Frenet trihedron of α . The coordinates in $\mathbf{T}$ and in $\mathbf{B}$ vanish. This means that

$$(1 + \lambda\kappa) \frac{\lambda\kappa'(1 + \lambda\kappa) - \lambda^2\tau\tau'}{[\lambda^2\tau^2 - (1 + \lambda\kappa)^2]^2} + \frac{\lambda\kappa'}{\lambda^2\tau^2 - (1 + \lambda\kappa)^2} = 0.$$

$$\lambda\tau \frac{\lambda\kappa'(1 + \lambda\kappa) - \lambda^2\tau\tau'}{[\lambda^2\tau^2 - (1 + \lambda\kappa)^2]^2} + \lambda\tau' \frac{\lambda\kappa'}{\lambda^2\tau^2 - (1 + \lambda\kappa)^2} = 0.$$

or

$$\lambda\tau(\lambda\kappa'\tau - \tau' - \lambda\kappa\tau') = 0.$$

$$\lambda(1 + \lambda\kappa)(\lambda\kappa'\tau - \tau' - \lambda\kappa\tau') = 0.$$

If $\tau = 0$ at some point, then $\tau \equiv 0$ and α would be a planar curve. If $\tau \neq 0$, then

$$\lambda\kappa'\tau - \tau' - \lambda\kappa\tau' = 0,$$

and this yields

$$\lambda\left(\frac{\kappa}{\tau}\right)' = -\left(\frac{1}{\tau}\right)'.$$

This implies that there is $b \in \mathbb{R}$ such that

$$\frac{\lambda\kappa}{\tau} + \frac{1}{\tau} = b.$$

Letting $A = -\lambda$ and $B = b$, we have proved a part of the theorem.

For the converse, that is, if there exist constants $A, B \in \mathbb{R}$ such that $A\kappa + B\tau = 1$, it suffices to define the curve

$$\beta(s) = \alpha(s) + A\mathbf{N}(s).$$

Then it is immediate that β is the mate curve of α . □

2.5 Exercises

1. Parametrize by the length-arc the curves that appeared at the beginning of this chapter.
2. Let α be a spacelike or timelike curve parametrized by the length-arc with non-zero acceleration. Show that α is a planar curve if and only if $\tau \equiv 0$.
3. Compute the Frenet equation of a planar spacelike curve whose vector normal is a lightlike vector.
4. Find an example of a curve α in a spacelike plane in such way that the curvature does not agree with the Euclidean curvature, when we see this curve as a map $\alpha : I \rightarrow \mathbf{E}^3$.
5. Study why is not possible extend Theorem 2.3.1 to the general case of planar curve, even in the case the this curve is spacelike.

Chapter 3

Spacelike and timelike surfaces in $\mathbf{E}_1^3$

In this chapter we introduce the notion of spacelike and timelike surfaces. We will define the mean curvature and the Gaussian curvature for this kind of surfaces. Next, we will locally write these curvatures and, finally, we will characterize umbilical surfaces of $\mathbf{E}_1^3$. The development of this chapter is similar to the Euclidean ambient, even in the local formulae of the curvature. However, we will see how the causal character imposes restrictions to work with these surfaces. For example, the surfaces can not be closed. Moreover, the Weingarten map for timelike surfaces is not diagonalizable (and we can not consider the notion of principal curvatures).

3.1 Surfaces in $\mathbf{E}_1^3$

Let M be a smooth, connected surface with boundary ∂M . Let $x : M \rightarrow \mathbf{E}_1^3$ be an *immersion*, that is, a differentiable map such that its differential map $dx_p : T_p M \rightarrow \mathbb{R}^3$ is injective. By the Inverse Theorem, x is a local homeomorphism on $x(M)$. If x is a global homeomorphism, we say that x is an *embedding* and that M is embedded (via x) in $\mathbf{E}_1^3$. If M is compact, this condition is equivalent to that $x(M)$ has not self-intersections.

We identify the tangent plane $T_p M$ with $(dx)_p(T_p M)$. We consider the pullback metric $g_p = x^*(\langle, \rangle_p)$, that is,

$$g_p(u, v) = \langle u, v \rangle_p = \langle dx_p(u), dx_p(v) \rangle.$$

In the Euclidean ambient $(T_p M, \langle, \rangle_p)$ is a Riemannian space, that is, the metric is positive definite. However, in the case that the map x arrives to $\mathbf{E}_1^3$, this metric can be of three types, namely,

1. $T_p M$ is a spacelike plane, that is, g_p is positive definite.
2. $T_p M$ is a timelike plane, that is, g_p is a metric with index 1.
3. $T_p M$ is a lightlike plane, that is, g_p is a degenerate metric.

Definition 3.1.1. *An immersion is called spacelike (resp. timelike, lightlike) if any tangent plane is spacelike (resp. timelike, lightlike).*

From now, we write $\langle, \rangle$ instead of g_p . Let us see that if the immersion is spacelike or timelike, we decompose the ambient space as $\mathbf{E}_1^3 = T_p M \oplus (T_p M)^\perp$, where the second subspace is a vector line, which will be timelike or spacelike if the immersion is spacelike or timelike, respectively.

The causal character of an immersion imposes conditions on the surface M . For example, we have:

Theorem 3.1.2. *Let $x : M \rightarrow \mathbf{E}_1^3$ be a spacelike immersion where M is a compact surface. Then $\partial M \neq \emptyset$.*

Proof: Consider the projection map $\pi(x, y, z) = (x, y)$. We define the map $X = \pi \circ x : M \rightarrow \mathbb{R}^2$. Assume that $\partial M = \emptyset$. As

$$|(dX)_p(u)|^2 = |(u_1, u_2, 0)|^2 = u_1^2 + u_2^2 > u_1^2 + u_2^2 - u_3^2 = |u|^2,$$

the map dX_p is an isomorphism and X is a local diffeomorphism. In particular, it is an open map and $x(M)$ is an open set of $\mathbb{R}^2$. On the other hand, M is compact and so, $X(M)$ is a closed set in $\mathbb{R}^2$. As conclusion, $X(M) = \mathbb{R}^2$: contradiction. $\square$

Therefore, if M is a compact surface, the boundary ∂M is a submanifold of dimension 1 and $M = \text{int}(M) \cup \partial M$. The tangent plane $T_p M$, with $p \in \partial M$, has bases where one of the vectors is tangent to ∂M and the other one is orthogonal to this one. This vector is called the unit conormal vector to M at p , and there exist interior and exterior conormal vector, depending if it points to $\text{int}(M)$ or not.

Corollary 3.1.3. *Let $x : M \rightarrow \mathbf{E}_1^3$ be a spacelike immersion of a compact surface. Assume that $x|_{\partial M}$ is a diffeomorphism between ∂M and a planar, closed, simple curve. Then $x(M)$ is a graph on the planar domain determined by $x(\partial M)$.*

Proof: Let Ω be the planar domain that encloses $x(\partial M)$. We know that $\pi : M \rightarrow \mathbb{R}^2$ is a local diffeomorphism. We claim that $\pi(M) \subset \Omega$. On the contrary, let $q \in \partial\pi(M) - \overline{\Omega}$ and let $x(p) = q$. Then $p \notin \partial M$ because $x(\partial M) = \partial\Omega$, and so, p is an interior point. But $x(p) \in \partial\pi(M)$ and then, the tangent plane at p must be vertical: contradiction. This shows the claim.

Thus $\pi \circ x : M \rightarrow \Omega$ is a local diffeomorphism. In particular, it is a covering map. Since Ω is simply connected, $\pi \circ x$ is a diffeomorphism. This means that $x(M)$ is a graph on Ω . $\square$

We now show examples of spacelike and timelike surfaces.

1. A horizontal plane is spacelike and a vertical plane is timelike.
2. The hyperboloid

$$\mathbb{H}^2 = \{p \in \mathbf{E}_1^3; \langle p, p \rangle = -1, z > 0\}$$

is a spacelike surface. For each $p \in \mathbb{H}^2$, the tangent plane is

$$T_p M = \{v \in \mathbb{R}^3; \langle p, v \rangle = 0\} = \langle p \rangle^\perp.$$

We call this surface *hyperbolic plane*. In section 3.4, we will justify its name. See figure 3.1.

3. The *De Sitter space* is defined as

$$\mathbb{S}_1^2 = \{p \in \mathbf{E}_1^3; \langle p, p \rangle = 1\}.$$

From an Euclidean viewpoint, this surface is a ruled hyperboloid. It is a timelike surface since $T_p M = \langle p \rangle^\perp$ and p is a spacelike vector. See figure 3.1.

4. The *lightlike cone*

$$\mathcal{C} = \{p \in \mathbf{E}_1^3; \langle p, p \rangle = 0\} - \{(0, 0, 0)\}$$

is a lightlike surface since $T_p M = \langle p \rangle^\perp$ and p is a lightlike vector.

5. Let f be a smooth map defined in a domain $\Omega \subset \mathbb{R}^2$ and let $S = \text{graph}(f)$. We consider the immersion $X : \Omega \rightarrow \mathbf{E}_1^3$ given by $X(x, y) = (x, y, f(x, y))$. As $X_x = (1, 0, f_x)$ and $X_y = (0, 1, f_y)$, the induced metric in the tangent plane has as a matrix

$$\begin{pmatrix} 1 - f_x^2 & -f_x f_y \\ -f_x f_y & 1 - f_y^2 \end{pmatrix}.$$

The determinant is

$$1 - f_x^2 - f_y^2 = 1 - |Df|^2.$$

Thus the immersion is spacelike if $|Df|^2 < 1$ and it is timelike if $|Df|^2 > 1$. We have the next examples:

- $f(x, y) = \sqrt{1 + x^2 + y^2}$ defined on $\Omega = \mathbb{R}^2$. This is the hyperbolic plane $\mathbb{H}^2$.
- $f(x, y) = \sqrt{1 + x^2}$. This surface is the ruled surface $z^2 - x^2 = 1$. It is a spacelike surface.




Figure 3.1: The hyperbolic plane, the De Sitter space and the lightlike cone.

Proposition 3.1.4. *Any spacelike surface is locally the graph of a function defined in the plane $\{z = 0\}$.*

Proof: We consider a local parametrization of the surface $X = X(u, v) = (x(u, v), y(u, v), z(u, v))$. Since the vector $X_u \times X_v$ is orthogonal to X_u and X_v , then it is a timelike vector. Thus, its third coordinate does not vanish. This coordinate is

$$-\begin{vmatrix} x_u & x_v \\ x_v & y_v \end{vmatrix}.$$

We use the Implicit Function Theorem to assert that around a point of the surface, the map

$$\phi : (u, v) \longmapsto (x(u, v), y(u, v))$$

is a diffeomorphism. We re-parametrize the immersion x by $X \circ \phi^{-1}$. Then it is immediate that it agrees with the graph of the function $z \circ \phi^{-1}$. $\square$

We end this section showing that any spacelike surface is orientable. Let us recall that a surface is orientable if there is a family of coordinate charts where the change of parameters have positive Jacobian. In Euclidean space we know that if x is an embedding with $\partial M = \emptyset$, then the surface is orientable. If the boundary is not empty, then there are examples of non-orientable surfaces, such as, the Moebius strip. If x is an immersion, there are examples of non-orientable surfaces in $\mathbf{E}^3$, as for example, the Klein bottle and the projective plane. If we delete a small piece to these surfaces, we obtain non-orientable immersions with boundary.

Theorem 3.1.5. *Let $x : M \rightarrow \mathbf{E}_1^3$ be a spacelike immersion. Then M is a orientable surface.*

Proof: Each tangent plane $T_p(M) = (dx)_p(T_p M)$ is spacelike. Locally we can define a unit vector N orthogonal to the tangent plane by the Lorentzian vector product. Exactly, let $x = x(u, v)$, $X_u = (dx)_p(\partial/\partial u)$ and $X_v = (dx)_p(\partial/\partial v)$. As the vectors X_u and X_v are spacelike, the vector product $X_u \times X_v$ is a timelike vector. We can define N as

$$N(u, v) = \frac{X_u \times X_v}{|X_u \times X_v|}.$$

Thus, in a neighbourhood of each point, there is a differentiable normal unit vector field on the surface.

Let $E_3 = (0, 0, 1)$. As this vector is also timelike, we have

$$|\langle N(p), E_3 \rangle| \geq 1.$$

Thus, either $\langle N(p), E_3 \rangle \geq 1$ or $\langle N(p), E_3 \rangle \leq -1 \forall p$ in a coordinate neighbourhood. For example, we choose that vector with negative sign. This allows us to define a globally normal vector field to the surface, that is, M is an orientable surface. $\square$

We now consider the map $N : M \rightarrow \mathbb{H}^2$ defined as above. We call N as the *Gauss map* of the immersion and it is globally defined. Moreover, it is future-directed. The choice of N gives us an orientation in $T_p M$: given a base $\{u, v\}$ of $T_p M$, we say that it is positive oriented if $\{u, v, N(p)\}$ is a positive oriented base of $\mathbf{E}_1^3$, that is, $\det(u, v, N(p)) > 0$.

The orientation on M defines an orientation on the boundary ∂M : given $u \in T_p \partial M$, we say that it is positive oriented if $\{u, v\}$ is a positive oriented base of $T_p M$, where v is the interior conormal vector.

The same argument shows that for timelike surfaces, there is defined locally a Gauss map.

3.2 Mean curvature of non-degenerate surfaces

We distinguish the cases that the surface is spacelike and timelike.

The spacelike case.

Let $x : M \rightarrow \mathbf{E}_1^3$ be a spacelike immersion of a surface M and let N be its Gauss map (pointing to the future). Let X be a vector field to M . We denote by ∇^0 and ∇ the Levi-Civita connections of $\mathbf{E}_1^3$ and M respectively. Moreover,

$$\nabla_X Y = (\nabla_X^0 Y)^\top$$

where the second term denotes the tangent part of the vector field $\nabla_X Y$. We define the second fundamental form of x as the tensorial, symmetric map $\sigma : \mathfrak{X}(M) \times \mathfrak{X}(M) \rightarrow \mathfrak{X}(M)^\perp$ by

$$\sigma(X, Y) = (\nabla_X^0 Y)^\perp.$$

The Gauss formula is

$$\nabla_X^0 Y = \nabla_X Y + \sigma(X, Y),$$

where X and Y are tangent vector fields to M . If Z is a normal vector field to x , we denote by $A_Z(X)$ the tangent component of $-\nabla_X^0 Z$, that is,

$$A_Z(X) = -(\nabla_X^0 Z)^\top.$$

We have

$$\langle A_Z(X), Y \rangle = \langle \sigma(X, Y), Z \rangle.$$

The map $A_Z : \mathfrak{X}(M) \rightarrow \mathfrak{X}(M)$ is called the Weingarten endomorphism associated to Z . Also, the map A_Z is linear and self-adjoint with respect to the metric of M . If we take $Z = N$, and because $\langle \nabla_X^0 N, N \rangle = 0$, we have $A_N(X) = -\nabla_X^0 N$. We say that $A := A_N$ is the *Weingarten endomorphism* of the immersion x . We have then $\langle AX, Y \rangle = \langle X, AY \rangle$. If $X, Y \in \mathfrak{X}(M)$,

$$\sigma(X, Y) = -\langle \sigma(X, Y), N \rangle N = -\langle A(X), Y \rangle N.$$

$$\nabla_X^0 Y = \nabla_X Y - \langle A(X), Y \rangle N.$$

As a consequence, the Weingarten endomorphism is diagonalizable, that is, if $p \in M$, the map $A_p : T_p M \rightarrow T_p M$ defined by $A_p(v) = (AX)_p$, where $X \in \mathfrak{X}(M)$ is a vector field that extends v , is diagonalizable. The eigenvalues of A_p are called *principal curvatures* and they will be denoted by $\lambda_i(p)$.

We define the *mean curvature vector field* of the immersion as

$$\vec{H} = \frac{1}{2} \text{traza}(\sigma).$$

Let $\{E_1, E_2\}$ be an orthonormal local vector fields. Then

$$\begin{aligned} \vec{H} &= \frac{1}{2} \text{traza}(\sigma) = \frac{1}{2} (\sigma(E_1, E_1) + \sigma(E_2, E_2)) = \\ &= -\frac{1}{2} (\langle AE_1, E_1 \rangle + \langle AE_2, E_2 \rangle) N = \left(-\frac{1}{2} \text{traza} A \right) N. \end{aligned}$$

If we want to have a identity $\vec{H} = HN$, we define the mean curvature of the immersion as the function

$$H = -\frac{1}{2} \text{traza} (A).$$

In particular, if $\vec{H} \neq 0$, then $\vec{H}$ is future-directed since $\langle \vec{H}, \vec{H} \rangle = -H^2 < 0$.

We compute the curvature tensor of M . As the curvature R^0 of $\mathbf{E}_1^3$ vanishes, the Gauss equation says

$$R(X, Y)Z = -\langle AX, Z \rangle AY + \langle AY, Z \rangle AX.$$

The Ricci curvature is

$$\text{Ric}(X, Y) = \sum_{i=1}^2 \langle R(X, E_i)Y, E_i \rangle = \langle AX, AY \rangle + 2H \langle AX, Y \rangle.$$

The scalar curvature is

$$\rho = \text{traza}(\text{Ric}) = -4H^2 + \text{trace}(A^2).$$

As $\rho = 2K$, where K is the *Gauss curvature*, then

$$K = -2H^2 + \frac{\text{traza}(A^2)}{2}.$$

The timelike case.

Let $x : M \rightarrow \mathbf{E}_1^3$ be a timelike immersion, that is, the induced metric on M via x is a non-degenerate metric of index 1. The surface is locally orientable: now N is a spacelike unit vector field and N is defined as $N : U \subset M \rightarrow \mathbb{S}_1^2$. Let us recall that there is not a Cauchy-Schwarz inequality for spacelike vectors.

Because it is possible, at least locally, to define N , we can do a similar development as the one for spacelike immersions. With the above notation, we have

$$\begin{aligned} \nabla_X^0 Y &= \nabla_X Y + \sigma(X, Y) = \nabla_X Y + \langle \sigma(X, Y), N \rangle N \\ &= \nabla_X Y + \langle AX, Y \rangle N. \end{aligned}$$

The Weingarten map A is self-adjoint again, that is,

$$\langle AX, Y \rangle = \langle X, AY \rangle \quad X, Y \in \mathfrak{X}(M).$$

The difference now is that A_p can be not diagonalizable¹.

However, we define the mean curvature and the Gauss curvature in the same way as the one for spacelike surfaces, that is,

$$H = \frac{1}{2} \text{traza}_I \sigma, \quad K = \det_I(\sigma),$$

where the subindex I denotes that we compute with respect to metric of the surface. For example, if $\{E_1, E_2\}$ is an orthonormal tangent vectors fields, with E_2 a timelike vector, then

$$H = \frac{1}{2} \left(\sigma(E_1, E_1) - \sigma(E_2, E_2) \right), \quad K = \sigma(E_1, E_1)\sigma(E_2, E_2) - \sigma(E_1, E_2)^2.$$

The De Sitter space $\mathbb{S}_1^2(r)$ is an example of a timelike surface. We compute the mean curvature and the Gauss curvature. Since $N(p) = p/r$, then $dN_p(v) = v/r$. This means that

$$\sigma_p(u, v) = \frac{1}{r} \langle u, v \rangle.$$

¹A self-adjoint endomorphism with respect to a metric that is not positive definite can be not diagonalizable. Exactly, the matrix of A_p with respect to an orthonormal base can be not symmetric.

Therefore

$$H \equiv \frac{1}{r}, \quad K \equiv \frac{1}{r^2}.$$

3.3 Local computations of curvatures

Let x be a spacelike immersion and let $X = X(u, v)$ be a local parametrization. Let $\{X_u, X_v\}$ be a local base of the tangent plane at each point. Let us recall that the first fundamental form is the metric on $T_p M$, that is,

$$I_p = \langle \cdot, \cdot \rangle_p : T_p M \times T_p M \rightarrow \mathbb{R}$$

$$I_p(u, v) = \langle u, v \rangle_p.$$

With respect to the base $B = \{X_u, X_v\}$, let $\begin{pmatrix} E & F \\ F & G \end{pmatrix}$ be the matricial expression of I , where

$$E = \langle X_u, X_u \rangle, \quad F = \langle X_u, X_v \rangle, \quad G = \langle X_v, X_v \rangle.$$

The metric is positive definite if and only if

$$\det(I) = EG - F^2 > 0.$$

If the immersion is timelike, then $EG - F^2 < 0$. We take the normal vector field given by

$$N = \frac{X_u \times X_v}{|X_u \times X_v|}.$$

The second fundamental form at p is

$$\sigma_p : T_p M \times T_p M \rightarrow \mathbb{R}$$

$$\sigma_p(u, v) = -\langle (dN)_p(u), v \rangle = \langle A_p(u), v \rangle.$$

Let $\begin{pmatrix} e & f \\ f & g \end{pmatrix}$ be the matricial expression of σ with respect to B , that is,

$$\begin{aligned} e &= -\langle X_u, N_u \rangle = \langle N, X_{uu} \rangle \\ f &= -\langle X_u, N_v \rangle = -\langle X_v, N_u \rangle = \langle N, X_{uv} \rangle \\ g &= -\langle X_v, N_v \rangle = \langle N, X_{vv} \rangle \end{aligned}$$

If $a, b \in T_p M$ (in coordinates with respect to B), we have

$$a^t \begin{pmatrix} e & f \\ f & g \end{pmatrix} b = a^t \begin{pmatrix} E & F \\ F & G \end{pmatrix} Ab.$$

Thus

$$A = \begin{pmatrix} E & F \\ F & G \end{pmatrix}^{-1} \begin{pmatrix} e & f \\ f & g \end{pmatrix}.$$

The mean curvature and the Gauss curvature are

$$\begin{aligned} H &= -\frac{1}{2} \frac{eG - 2fF + gE}{EG - F^2} \\ K &= -\frac{eg - f^2}{EG - F^2} \end{aligned}$$

In the case that the surface is timelike, the formulas only change of sign.

In order to show the difference between the spacelike and timelike case, we are going to study a timelike surface whose Weingarten endomorphism is not diagonalizable. The surface is a ruled surface. Let $\alpha : I \rightarrow \mathbf{E}_1^3$ be a lightlike curve and we denote by $\{\mathbf{T}, \mathbf{N}, \mathbf{B}\}$ the Frenet trihedron. The binormal vector is a unit spacelike vector. We consider the map $X : I \times \mathbb{R} \rightarrow \mathbf{E}_1^3$ given by

$$X(s, t) = \alpha(s) + t\mathbf{B}(s).$$

We compute the matrix of A_p with respect to the metric $\{X_s, X_t\}$. Since $X_s = \mathbf{T} + t\mathbf{B}' = \mathbf{T} - t\tau\mathbf{N}$ and $X_t = \mathbf{B}$, then

$$\begin{pmatrix} t^2\tau^2 & 1 \\ 1 & 0 \end{pmatrix}.$$

As the determinant is negative, the surface is timelike. We compute the coefficients of the second fundamental form. For this, we have

$$X_{ss} = (1 - t\tau^2)\mathbf{T} + (1 - t\tau')\mathbf{N} + t\tau\mathbf{B}, \quad X_{st} = -\tau\mathbf{N}, \quad X_{tt} = 0.$$

Then the second fundamental form is

$$\begin{pmatrix} -1 + t(-\tau + \tau' + t^2\tau^3) & \tau \\ \tau & 0 \end{pmatrix}.$$

The Weingarten endomorphism is now

$$A = \begin{pmatrix} \tau & 0 \\ -1 + t\tau(-1 + (-1 + t)t\tau^2) + t\tau' & \tau \end{pmatrix}.$$

This matrix is not diagonalizable. On the other hand, the mean curvature is $H = \tau$ and the Gauss curvature is $K = \tau^2$. Finally, let us see that this surface satisfies $H^2 = K$ but it is not umbilical.

3.4 Umbilical surfaces

In this section, we find all (spacelike or timelike) umbilical surfaces of $\mathbf{E}_1^3$. Let $x : M \rightarrow \mathbf{E}_1^3$ be a spacelike or timelike immersion.

Definition 3.4.1. A point $p \in M$ is called umbilical if

$$\sigma_p(u, v) = \lambda(p)\langle u, v \rangle,$$

that is, the second fundamental form is proportional to the metric. In the case that the immersion is spacelike, this is equivalent to say that $\lambda_1(p) = \lambda_2(p)$. A surface is called totally umbilical if any point is umbilical.

Let x be a spacelike immersion. As

$$H = -\frac{\lambda_1 + \lambda_2}{2}, \quad K = -\lambda_1 \lambda_2,$$

where λ_i are the eigenvalues of A . Then

$$\left(\frac{\lambda_1 - \lambda_2}{2}\right)^2 = \left(\frac{\lambda_1 \lambda_2}{2}\right)^2 - \lambda_1 \lambda_2 = H^2 + K.$$

Thus in a spacelike surface it holds $H^2 + K \geq 0$ and it is zero if and only if the point is umbilical. On the other hand, the principal curvatures are

$$\lambda_i = -H \pm \sqrt{H^2 + K}.$$

We consider the next examples of umbilical surfaces:

1. A spacelike or timelike plane $P = p_0 + \langle v \rangle^\perp$, $|v| = 1$, satisfies $N = v$. Then $dN = 0$, and the surface is totally umbilical with $\lambda_i \equiv 0$ in the case of spacelike planes.
2. Let $\mathbb{H}^2(r, p_0)$ be the hyperbolic plane

$$\mathbb{H}^2(r, p_0) = \{p \in \mathbf{E}_1^3; \langle p - p_0, p - p_0 \rangle = -r^2, z > 0\}.$$

Then $N(p) = p/r$ and $dN_p(v) = v/r$. Then the surface is totally umbilical with $\lambda_i \equiv -1/r$. Here $H = 1/r$ and $K = -1/r^2$.

We justify the name of hyperbolic plane to the surface $\mathbb{H}^2$. For each point $p \in \mathbb{H}^2$ and $v \in T_p \mathbb{H}^2$ with $|v| = 1$, we define the curve $\alpha(s) = \cosh(s)p + \sinh(s)v$. Then $\alpha(s) \in \mathbb{H}^2$, with $\alpha(0) = p$ and $\alpha'(0) = v$. Because $\alpha''(s) = \alpha(s)$, the tangent part of the acceleration vanishes, that is, $\alpha''(s)^T = 0$. This means that α is a geodesic. Because the tangent vector v is arbitrary, all geodesic starting from p are written as above. As α is defined for any $s \in \mathbb{R}$, the surface is complete. Then $\mathbb{H}^2$ is a complete, simply connected surface with constant curvature $K \equiv -1$. A classical result in Differential Geometry assures that $\mathbb{H}^2$ is isometric to the hyperbolic plane.

3. Let $\mathbb{S}_1^2(r, p_0)$ be the De Sitter space

$$\mathbb{S}_1^2(r, p_0) = \{p \in \mathbf{E}_1^3; \langle p - p_0, p - p_0 \rangle = r^2\}.$$

Then $N(p) = p/r$ and $dN_p(v) = v/r$ and so, the surface is totally umbilical. Here $H \equiv 1/r$ and $K \equiv 1/r^2$.

Theorem 3.4.2. *The only totally umbilical surfaces in Minkowski space are planes, hyperbolic planes or De Sitter spaces.*

Proof: We work in a coordinate neighbourhood and let $X = X(u, v)$ be the corresponding parametrization. As the surface is totally umbilical, there is a smooth function f such that

$$\begin{aligned} (N \circ X)_u &= (f \circ X)X_u \\ (N \circ X)_v &= (f \circ X)X_v \end{aligned}$$

A differentiation with respect to u and v yields

$$(f \circ X)_u X_v + (f \circ X) X_{uv} = (f \circ X)_v X_u + (f \circ X) X_{uv}.$$

Thus $(f \circ X)_u = (f \circ X)_v$. This means that f is a constant function in a coordinate neighbourhood, that is, $f \circ X = r$, $r \in \mathbb{R}$. Since the surface is connected, $f \circ X = r$ on M .

1. If $r = 0$, then $N_u = N_v$, that is, N is constant. This shows that the surface is a plane.
2. If $r \neq 0$, then $N_u = rX_u$ and $N_v = rX_v$. We define in a neighbourhood the function

$$h(u, v) = X(u, v) - \frac{1}{r}(N \circ X)(u, v).$$

Then $h_u = h_v = 0$. This means that h is constant. Thus there exists $p_0 \in \mathbf{E}_1^3$ such that

$$X(u, v) - \frac{1}{r}(N \circ X)(u, v) = p_0.$$

Then

$$\langle X - p_0, X - p_0 \rangle = \mp \frac{1}{r^2},$$

with the sign depending if the surface is spacelike or timelike, respectively. Anyway, the surface is a hyperbolic plane or De Sitter space.

□

3.5 Exercises

1. Show that any timelike surface is locally a graph on the plane $\{y = 0\}$ or the plane $\{x = 0\}$.
2. Let M be a spacelike surface in $\mathbf{E}_1^3$ and $p \in M$. Show that if $K(p) < 0$, there exists a neighbourhood of p where the surface lies in one side of $T_p M$. If $K(p) > 0$, then in any neighbourhood of p there are points in both sides of $T_p M$.
3. Compute K and H for the next surfaces
 - $x^2 + y^2 = 1$.
 - $x^2 - y^2 = 1$.
 - $z^2 - x^2 = 1$.

Chapter 4

Spacelike surfaces with constant mean curvature

We characterize constant mean curvature (CMC) spacelike surfaces as solutions of a variational problem. We state the maximum principle and we compute the Laplacian of two functions. The statements of results are similar as in Euclidean space. However, and as we will see in the next chapter, the consequences are strong different.

4.1 The variational problem

The results that we are going to obtain are similar than in Euclidean space with minor modifications.

Let M be a surface and let $x : M \rightarrow \mathbf{E}_1^3$ be a spacelike immersion in $\mathbf{E}_1^3$. A *variation* of x is a differentiable map $X : (-\epsilon, \epsilon) \times M \rightarrow \mathbf{E}_1^3$ that satisfies the following properties (see figure 4.1):


1. $X(0, p) = x(p)$, $\forall p \in M$.
2. The maps $X_t : M \rightarrow \mathbf{E}_1^3$ given by $X_t(p) = X(t, p)$ are spacelike immersions for each $t \in (-\epsilon, \epsilon)$.
3. If M is a compact surface ($\partial M \neq \emptyset$), we suppose that $X(t, p) = x(p)$ for $p \in \partial M$. This condition means that the immersions of the variation leave pointwise fixed the boundary of M .

We define the variational field throughout X as

$$\frac{\partial X}{\partial t}(t, p) := (dX)_{(t, p)}\left(\frac{\partial}{\partial t}, 0\right).$$

At $t = 0$, we define *variational vector field* of the variation X as

$$\xi(p) = \frac{\partial X}{\partial t}(0, p).$$

Figure 4.1: A variation of an immersion x .

The variational vector field is called *normal* if it is orthogonal to each tangent plane, that is, $\xi(p) \perp T_p M$, $p \in M$. Thus we write

$$\xi(p) = f(p)N(p)$$

for some smooth function f in M .

From now, we suppose that M is a compact surface. We define the *area functional* as

$$A(t) = \int_M dA_t,$$

where dA_t is the area element on M induced by the metric $X_t^*(\langle, \rangle)$. At $t = 0$, the value $A(0)$ agrees with the area of $x(M)$.

We define the *volume functional* as

$$V(t) = \frac{1}{3} \int_M \langle X_t, N_t \rangle dA_t,$$

where N_t is a unit normal vector field to the immersion X_t . The value $V(t)$ represents the enclosed volume by the cone whose base is $X_t(M)$ and vertex the origin of coordinates. In Euclidean ambient and in the case that the surface is closed, $V(t)$ is the volume enclosed by the domain $X_t(M)$.

Because $\mathbf{E}_1^3 = T_p M \oplus T_p M^\perp$, we decompose the variational vector field ξ as

$$\xi(p) = \xi(p)^T + f(p)N(p), \quad f \in C^\infty(M).$$

For the tangent part $\xi(p)^T$, there exists $\xi' \in \mathfrak{X}(M)$ such that $(dx)_p \xi'(p) = \xi(p)^T$. The map $A(t)$ is differentiable at $t = 0$. Moreover

$$A'(0) = -2 \int_M (Hf) dA - \int_{\partial M} \langle \xi', \nu \rangle ds,$$

where H is the mean curvature of the immersion and ν is the unit conormal vector field along ∂M . Since the variation X fixes the boundary ∂M , the second summand vanishes (if $p \in \partial M$, the map $t \mapsto X(t, p)$ is constant, and then, $t \mapsto \frac{\partial X}{\partial t}(t, p)$ is 0).

Theorem 4.1.1. *Let $x : M \rightarrow \mathbf{E}_1^3$ be a spacelike immersion and let X be a variation of x . The first variation of the area is*

$$A'(0) = 2 \int_M (Hf) dA. \quad (4.1)$$

For the volume functional and for variations that preserve the boundary, we have the next formula

$$V'(0) = - \int_M f \, dA. \quad (4.2)$$

If the variation preserves the volume, then the function $V(t)$ is constant and so, $V'(0) = 0$. Then $\int_M f \, dA = 0$. Conversely, one can prove that if $f \in C^\infty(M)$ satisfies $\int_M f \, dA = 0$, then there is a variation X that fixes the boundary and whose variational vector field is fN .

Theorem 4.1.2. *Let M be a compact surface and let $x : M \rightarrow \mathbf{E}_1^3$ be a spacelike immersion in $\mathbf{E}_1^3$. Then x has constant mean curvature if and only if $t = 0$ is a critical point of the area functional for any volume preserving variation of x that fixes the boundary.*

Proof: Suppose that x has constant mean curvature H . Let X be a variation of x that preserves the boundary and the volume of x . Then $V'(0) = 0$ and so, $\int_M f \, dA = 0$. Since H is constant, the equality (4.1) writes as

$$A'(0) = 2 \int_M Hf \, dA = 2H \int_M f \, dA = 0.$$

For the converse, let H be the mean curvature of x and let $c \in \mathbb{R}$ be the number defined by

$$c = \frac{\int_M H \, dA}{A(M)}, \quad A(M) := \text{area}(x(M)).$$

We consider the function $f = H - c$. Since $\int_M f \, dA = 0$, we know that there exists a volume preserving variation that fixes the boundary and with variational vector field fN . From (4.1) we have

$$0 = A'(0) = 2 \int_M (H - c)f \, dA = 2 \int_M (H - c)^2 \, dA.$$

Thus $H - c \equiv 0$, that is, H is a constant function. $\square$

It is possible to do a new variational characterization for CMC surfaces with variations that do not fix the volume of the immersion. Exactly, let X be a variation of a spacelike immersion x that fixes the boundary of M . Given $c \in \mathbb{R}$, we define the functional J_c by

$$J_c(t) = A(t) + 2c V(t).$$

Then

$$J'_c(0) = A'(0) + 2cV'(0) = 2 \int_M (H - c)f \, dA,$$

where $\xi^\perp = fN$.

Theorem 4.1.3. *A spacelike surface has constant mean curvature if and only if there exists $c \in \mathbb{R}$ such that $J'_c(0) = 0$ for any variation that fixes the boundary.*

Proof: If H is constant, then $J'_H(0) = 0$. Assume now that there is a number c such that $J'_c(0) = 0$. We use Theorem 4.1.2: let X be a variation preserving the volume and the boundary. Because $V'(0) = 0$, then $0 = J'_c(0) = A'(0)$. $\square$

4.2 The maximum principle

In this section we state the maximum principle for spacelike surfaces with constant mean curvature. The maximum principle is a consequence from a result of partial differential equations theory, named the maximum principle again. Because its fundament is part of this theory, we are not going to do a proof of the result: our setting is a particular case. Moreover, the next presentation holds in Lorentzian as well as Euclidean ambient. The maximum principle is used to "compare" two surfaces that have the *same* mean curvature.

Let S_1 and S_2 be two surfaces that are tangent at a common point $p \in S_1 \cap S_2$. Suppose that one of the surfaces, for example S_1 , lies below the other one around the point p . Exactly we consider both surfaces as graphs of smooth functions u_1 and u_2 on a domain of the common tangent plane $T_p S_1 = T_p S_2$. After an isometry, we assume that $T_p S_i$ is the horizontal plane $z = 0$ (here we use the spacelike condition). We take orientation in both graphs in such way that they agree at p , that is, $N_1(p) = N_2(p)$. In such situation, we say that S_1 lies below S_2 if $u_1 \leq u_2$ in a neighbourhood of p in the tangent plane. We write then $S_1 \leq S_2$ (or $S_2 \geq S_1$).

Even in the case that p is a boundary point, we have the next definition (see figure 4.2)

Definition 4.2.1. *Let S_1 and S_2 be two spacelike surfaces in $\mathbf{E}_1^3$, with $p \in S_1 \cap S_2$ and such that $N_1(p) = N_2(p)$. We say that $S_1 \leq S_2$ around p if, when we write the surfaces as graphs on the common tangent plane, $S_i = \text{graph}(u_i)$, we have $u_1 \leq u_2$ in a neighbourhood of p . If $p \in \partial S_1 \cap \partial S_2$, we add the condition $T_p \partial S_1 = T_p \partial S_2$.*




Figure 4.2: Tangent point between two surfaces, with $S_1 \leq S_2$.

It is easy to show that if $u_1 \leq u_2$, then $H_1(p) \leq H_2(p)$ ¹. In the literature this result is called sometimes the comparison principle.

As an example to apply the comparison principle is the following. Let S be a surface with constant mean curvature $H \neq 0$ such it appears in the figure 4.3. We want to know if with the orientation pointing upwards, the mean curvature is positive or it is negative. In the lowest point of S , we take the tangent plane P . Then $P \leq S$. The plane P has zero mean curvature (independently of the orientation). On P we take the orientation pointing upwards. Then $P \leq S$ and the comparison principle yields $0 \leq H(p)$. As $H \neq 0$, then $H(p)$ is positive, and as H is constant, $H > 0$ in S .

¹After a change of coordinate, we suppose that $p = (0, 0, 0)$. This means that $\partial u_i / \partial x$ and $\partial u_i / \partial y$ vanish at p . If we define the function $f = u_2 - u_1$, the point p is a minimum of f ; in particular, the hessian $D^2 f(p)$ is positive semi-definite and its trace satisfies $2(H_2(p) - H_1(p)) \geq 0$




Figure 4.3: The comparison principle

The maximum principle refers to the fact that if two surfaces with the same mean curvature, one of them lies below the other one, then both surfaces agree in an *open set* around p .

Theorem 4.2.2 (Maximum principle). *Let S_1 and S_2 be two spacelike surfaces with a common (interior or boundary) tangent point p . Suppose that $S_1 \leq S_2$. If the mean curvatures agree and are constant, then $S_1 = S_2$ in a neighbourhood of p . See figure 4.4.*




Figure 4.4: The maximum principle.

We are going to indicate how the proof works (the same argument holds for CMC surfaces in $\mathbf{E}^3$). Let S be a graph of a function u , that is, $S = \{(x, y, u(x, y)); (x, y) \in \Omega\}$, where $u \in C^\infty(\Omega)$ and Ω is a domain of $\mathbb{R}^2$. Consider in S the induced metric from $\mathbf{E}_1^3$ or $\mathbf{E}^3$. In the first case, if the surface is spacelike, the function u satisfies $|Du| < 1$ in Ω . We consider the orientation N that points upwards. Then N writes as

$$N = \frac{(-\epsilon Du, 1)}{\sqrt{1 + \epsilon |Du|^2}},$$

where $\epsilon = 1$ if the surface lies in $\mathbf{E}^3$ and $\epsilon = -1$ if it is in $\mathbf{E}_1^3$. Then the mean curvature H is

$$H = \epsilon \frac{1}{2} \operatorname{div} \left(\frac{Du}{\sqrt{1 + \epsilon |Du|^2}} \right).$$

This equality writes as

$$(1 + \epsilon u_y^2) - 2\epsilon u_x u_y u_{xy} + (1 + \epsilon u_x^2) u_{yy} = 2H\epsilon(1 + \epsilon(u_x^2 + u_y^2))^{3/2}.$$

This is a partial differential equation, and we put

$$Qu := \sum_{ij} a_{ij}(x, y, u_x, u_y) D_{ij}u + b(x, y, u_x, u_y) \quad D_{ij}u = \frac{\partial^2 u}{\partial x_i \partial x_j}.$$

From the PDE theory and in order to classify Q , it is necessary to study the matrix $A = (a_{ij})$ of the coefficients of second order. In the above case, such matrix is

$$\begin{pmatrix} 1 + \epsilon u_y^2 & -\epsilon u_x u_y \\ -\epsilon u_x u_y & 1 - \epsilon u_x^2 \end{pmatrix}.$$

This matrix is positive definite because

$$\det \begin{pmatrix} 1 + \epsilon u_y^2 & -\epsilon u_x u_y \\ -\epsilon u_x u_y & 1 - \epsilon u_x^2 \end{pmatrix} = 1 + \epsilon |Du|^2 > 0.$$

This means that $Qu = 0$ is an elliptic equation. See that this is immediate if $\epsilon = 1$, but if $\epsilon = -1$, the spacelike condition is basic. *Linear* elliptic equations have the property that satisfy a maximum principle in the sense that if two functions satisfy the same equation (and the same boundary conditions), then both solutions agree.

In our case, Q is not linear on the variables Du_{ij} . Exactly, an equation of type $Qu = 0$ is called a *quasilinear equation*. The reason is the following: if u_1 and u_2 satisfy the same equation $Qu_i = 0$, the difference function $u = u_1 - u_2$ satisfies a linear elliptic operator, that is,

$$Lu := Q(u_1) - Q(u_2),$$

and L is a linear operator. Then we can apply the maximum principle to the function u . This proves Theorem 4.2.2

.

4.3 Two equations for CMC spacelike surfaces

In this section we compute the Laplacian of two functions associated to a spacelike immersion. Let $x : M \rightarrow \mathbf{E}_1^3$ be a spacelike immersion and let $a \in \mathbf{E}_1^3$. We are going to compute $\Delta \langle x, a \rangle$ and $\Delta \langle N, a \rangle$, where Δ is the Laplace operator in $(M, \langle, \rangle)$. The Laplacian of a function $f \in C^\infty(M)$ is defined as

$$\Delta f = \operatorname{div} \nabla f = \operatorname{traza} \left(v \longmapsto \nabla_v \nabla f \right).$$

We compute $\Delta f(p)$, $p \in M$ by taking an adapted orthonormal base at p , that is, an orthonormal base of tangent vector fields $\{E_1, E_2\}$ such that $\nabla_{E_i(p)} E_j = 0$, $i, j \in \{1, 2\}$. In order to emphasize that the computations are done at p , we indicate $e_i = E_i(p)$. With respect to this base, the Laplacian is

$$\Delta f(p) = \sum_{i=1}^2 e_i(E_i(f)).$$

In the case of $\Delta \langle N, a \rangle$ we suppose that the mean curvature is constant.

1. Let $f = \langle x, a \rangle$. Then $E_i(f) = \langle E_i, a \rangle$ and at the point p we have

$$\begin{aligned} e_i(E_i(f)) &= e_i \langle E_i, a \rangle = \langle \nabla_{e_i}^0 E_i, a \rangle = \langle \nabla_{e_i} E_j + \sigma(e_i, e_j), a \rangle \\ &= \langle \sigma(e_i, e_j), a \rangle = -\langle Ae_i, e_i \rangle \langle N, a \rangle. \end{aligned}$$

Thus

$$\Delta \langle x, a \rangle = - \left(\sum_{i=1}^2 \langle Ae_i, e_i \rangle \right) \langle N, a \rangle = 2H \langle N, a \rangle.$$

2. Suppose now that H is constant. First we show that $\sum_{i=1}^2 \nabla_{e_i}^0 \nabla_{E_i}^0 N$ is orthogonal to the surface, that is,

$$\sum_{i=1}^2 \langle \nabla_{e_i}^0 \nabla_{E_i}^0 N, e_k \rangle = 0, \quad k = 1, 2. \quad (4.3)$$

Since $\langle N, E_k \rangle = 0$, then $\langle \nabla_{E_i}^0 N, E_k \rangle = -\langle N, \nabla_{e_i}^0 E_k \rangle$. Therefore

$$\langle \nabla_{e_i}^0 \nabla_{E_i}^0 N, e_k \rangle + \langle \nabla_{e_i}^0 N, \nabla_{e_i}^0 E_k \rangle = -\langle \nabla_{e_i}^0 N, \nabla_{e_i}^0 E_k \rangle - \langle N, \nabla_{e_i}^0 \nabla_{e_i}^0 E_k \rangle.$$

As $\nabla_{e_i}^0 N$ is tangent and $\nabla_{e_i} E_k = 0$, then

$$\langle \nabla_{e_i}^0 \nabla_{E_i}^0 N, e_k \rangle = -\langle N, \nabla_{e_i}^0 \nabla_{e_i}^0 E_k \rangle = -\langle N, \nabla_{e_i}^0 \nabla_{E_k}^0 E_i \rangle.$$

The metric in $\mathbf{E}_1^3$ is flat and this means $\nabla_{e_i}^0 \nabla_{E_k}^0 E_i = \nabla_{e_k}^0 \nabla_{E_i}^0 E_i$. By substituting in the above equation, we obtain:

$$\langle \nabla_{e_i}^0 \nabla_{E_i}^0 N, e_k \rangle = -\langle \nabla_{e_k}^0 \nabla_{E_i}^0 E_i, N \rangle,$$

and

$$\sum_{i=1}^2 \langle \nabla_{e_i}^0 \nabla_{E_i}^0 N, e_k \rangle = -\langle \nabla_{e_k}^0 \left(\sum_{i=1}^2 \nabla_{E_i}^0 E_i \right), N \rangle. \quad (4.4)$$

The same computation holds in Euclidean space. As above, we use ϵ to distinguish the two ambient spaces. The mean curvature is $2H = \epsilon \sum_{i=1}^2 \langle \nabla_{e_i}^0 E_i, N \rangle$. Because H is constant,

$$e_k \left\langle \sum_{i=1}^2 \nabla_{e_i}^0 E_i, N \right\rangle = 0,$$

that is,

$$\langle \nabla_{e_k}^0 \left(\sum_{i=1}^2 \nabla_{E_i}^0 E_i \right), N \rangle + \left\langle \sum_{i=1}^2 \nabla_{E_i}^0 E_i, \nabla_{e_k}^0 N \right\rangle = 0.$$

The second summand vanishes since that the left part is orthogonal to the surface and the right one is tangent. From (4.4),

$$\langle \nabla_{e_k}^0 \left(\sum_{i=1}^2 \nabla_{E_i}^0 E_i \right), N \rangle = 0,$$

and we have proved the claim.

On the other hand,

$$\sum_{i=1}^2 \langle \nabla_{e_i}^0 \nabla_{E_i}^0 N, N \rangle = - \sum_{i=1}^2 \langle \nabla_{e_i}^0 N, \nabla_{e_i}^0 N \rangle = - \sum_{i=1}^2 \langle Ae_i, Ae_i \rangle \quad (4.5)$$

$$= - \sum_{i=1}^2 \langle A^2 e_i, e_i \rangle = -\text{trace } (A^2). \quad (4.6)$$

From (4.3) and (4.6), we conclude

$$\Delta \langle N, a \rangle = \sum_{i=1}^2 E_i(E_i \langle N, a \rangle) = -\epsilon \text{ traza } (A^2) \langle N, a \rangle.$$

We summarize the above computations in the next

Theorem 4.3.1. *Let $x : M \rightarrow \mathbf{E}_1^3$ be a spacelike immersion and let N be an orientation on M . Given $a \in \mathbf{E}_1^3$, we have*

$$\Delta \langle x, a \rangle = 2H \langle N, a \rangle.$$

Here H is the mean curvature of the immersion. Furthermore, if H is constant, then

$$\Delta \langle N, a \rangle = \text{trace}(A^2) \langle N, a \rangle.$$

Finally, recall that

$$\text{trace } (A^2) = 4H^2 + 2K,$$

and that $H^2 + K \geq 0$ in any point of the surface. Moreover $H^2 + K = 0$ only at the umbilical points. Thus, the last equation writes as

$$\Delta \langle N, a \rangle - (4H^2 + 2K) \langle N, a \rangle = 0.$$

In the case that the immersion arrives into the Euclidean space $\mathbf{E}^3$, the last equation is $\Delta \langle N, a \rangle = -\text{traza}(A^2) \langle N, a \rangle$, o $\Delta \langle N, a \rangle + (4H^2 - 2K) \langle N, a \rangle = 0$. As consequence, we write the equation in both ambients as

$$\Delta \langle N, a \rangle + \epsilon(4H^2 - \epsilon 2K) \langle N, a \rangle = 0.$$

Chapter 5

Compact surfaces with constant mean curvature

The focus of this chapter is the family of compact surfaces with constant mean curvature. In particular, the boundary is a non-empty set. If $C \subset \mathbf{E}_1^3$ is a curve in the ambient space, we say that an immersion $x : M \rightarrow \mathbf{E}_1^3$ has boundary C if the restriction map $x|_{\partial M}$ is a diffeomorphism between ∂M and C .

It is important to recall that if C is a simple, planar, closed curve, then x is a graph on the enclosed domain by C .

This chapter has two parts. The first one obtains consequence of the maximum principle for this kind of surfaces. The second one refers to the Dirichlet problem for the constant mean curvature equation. In both cases, the results obtained are different depending on the ambient space. In this sense, it is a bit surprise because the theoretical framework is the same for both kind of surfaces.

5.1 Consequences of the maximum principle

In this section, we consider a spacelike immersion $x : M \rightarrow \mathbf{E}_1^3$ with constant mean curvature, where M is a compact surface and $C \subset \mathbf{E}_1^3$ is the boundary of the immersion. We say M , x or $x(M)$. We denote by $\mathbb{H}^2(r)$ the hyperbolic plane of radius r and orientation according to the future direction. We know that its mean curvature is $H = 1/r$.

Theorem 5.1.1. *If M is a compact CMC spacelike surface with $H \neq 0$ and boundary included in a plane P , then the surface lies in one side of P .*

Proof: Without loss of generality, let P be a horizontal plane and assume that $H > 0$. We are going to show that the surface lies below P . By contradiction, we assume that there exist points above P . We take a horizontal plane in the highest point and tangent to the surface. The comparison principle gets a contradiction. $\square$

Let us see that the boundary of M can have some connected components and that M can be not a

graph. With the same reasoning, and assuming that the boundary is not a planar curve, we have

Corollary 5.1.2. *Let M be a spacelike surface with $H \equiv 0$ and with boundary $\partial S = C$. Then M lies included in the spacelike convex hull of C .*

Proof: It suffices to compare the surface with planes. Because there are not interior contact points (both surfaces have the same mean curvature $H = 0$ with any orientation), the plane can move until to arrive the boundary. We point out that the spacelike convex hull of C is the convex hull but using only spacelike planes. $\square$

The next result says us that two graphs with the same mean curvature and the same boundary agree. This is a consequence of the maximum principle of a solution of a quasilinear elliptic equation. However, we are going to do a "geometric" proof of this result.

Theorem 5.1.3. *Let C be a curve in $\mathbf{E}_1^3$ (not necessarily a planar curve). If S_1 and S_2 are two graphs with the same mean curvature and the same boundary C , then $S_1 = S_2$.*

Proof: We suppose that the surfaces are graphs on a horizontal plane P . Let S_1 and we move it vertically upwards until it does not touch S_2 (this is possible because both surfaces are compact). Next, we descend S_1 until the first time that it touches S_2 . If there is an interior tangent point $p \in S_1 \cap S_2$, this can not occur before S_1 returns to its original position. Because both surfaces have the same mean curvature and $S_1 \geq S_2$, the maximum principle says that both surfaces agree around p . By connection, $S_1 = S_2$.

If the contact point is a boundary point, this means that S_1 has returned its original position and $S_2 \leq S_1$. There are two possibilities:

1. If the contact point is a tangent point, the maximum principle concludes that $S_1 = S_2$ again.
2. On the contrary, the slope of S_1 along C is strictly bigger than the one of S_2 . Now we descend S_1 until it does not touch S_2 . Next, we move it upwards. Since in its original position, S_1 lies strictly above S_2 along C , then there exists an interior tangent point between S_1 and S_2 at some time t before the original position. If we denote by $S_1(t)$ the surface S_1 at this position, the maximum principle yields $S_1(t) = S_2$. But this is impossible because $\partial S_1(t) \neq C = \partial S_2$. This contradiction says that this situation can not occur.

$\square$

For the next result, we need to introduce the concept of hyperbolic cap. If $R, r > 0$, a *hyperbolic cap* is defined by

$$C(r; R) = \{p \in \mathbf{E}_1^3; x^2 + y^2 - z^2 = -r^2, z > 0, z^2 \leq r^2 + R^2\} \subset \mathbb{H}^2(r).$$

This surface is spacelike with mean curvature $H = 1/r$ with the upwards orientation, that is, $N(p) = p/r$. The boundary of $C(r; R)$

$$\partial C(r; R) = \{p \in C(r; R); z^2 = r^2 + R^2\}$$

is a circle in the spacelike plane $\{z = \sqrt{R^2 + r^2}\}$.

Corollary 5.1.4. *Let C be a circle in $\mathbf{E}_1^3$, that is, a planar, closed spacelike curve with constant curvature. Then the only spacelike CMC surfaces with boundary C are planar discs and hyperbolic caps.*

Proof: Because C is a simple closed curve, the surface must be a graph. After an isometry, we assume that the plane containing C is a horizontal plane. Let S be a graph with mean curvature H and $\partial S = C$. Then S is a graph on the domain bounded by C . If $H = 0$, the planar disc that bounds C is a graph with $H = 0$. By the uniqueness of Theorem 5.1.3, S is a planar disc.

Let $H \neq 0$. We have only to show that there exists a hyperbolic cap with mean curvature H and boundary C . Assume that the radius of C is R . Then the cap that we are looking for is $C(\frac{1}{H}; R)$. $\square$

Remark that the above result is not true in the Euclidean ambient, that is, there are compact CMC surfaces with circular boundary that are not umbilical. These surfaces are not embedded with higher genus. However, it is an open problem to know if planar discs and spherical caps are the only CMC surfaces with circular boundary that are embedded or are topological discs.

The above result can be proved using the Alexandrov reflection method¹.

Theorem 5.1.5. *Let S be a compact CMC surface with boundary C included in a plane P . If S makes constant angle with P along C then S is a domain of P or C is a circle and S is a hyperbolic cap.*

Let us see that ∂S can have some connected component and that S is not necessarily a graph.

Proof: Let denote by Ω the domain bounded by C in P . If $H = 0$, the comparison principle asserts that S can not have points in both sides of P , that is, $S = \Omega$.

Suppose that $H \neq 0$. After an isometry, we take P as a horizontal plane. We know that S lies in one side of P ; exactly, if $H > 0$, S lies below P . If $C = C_1 \cup \dots \cup C_k$ is the decomposition into connected components of C , we attach to S appropriate domains Ω_i to get that $S \cup (\Omega_1 \cup \dots \cup \Omega_n)$ is a closed (non-smooth) surface M . Let W be the domain that encloses M . See figure 5.1.




Figure 5.1: The surface M and the domain W .

Let v be a horizontal direction. We take the family of orthogonal parallel planes to v , that is, it is a family of vertical, parallel planes. The symmetry with respect to these planes are isometries of $\mathbf{E}_1^3$, because the corresponding matrix with respect to the usual base is

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

We suppose that $v = (0, 1, 0)$, so we can talk of "right" and "left". We take a plane of the family far away from S and on the right of S . By moving on the left in the direction given by v we arrive

¹Alexandrov showed with this technique that the only closed embedded CMC surfaces of $\mathbf{E}^3$ are round spheres.

until the first contact point with M . After this moment, we move a little on the left and we reflect the right side of S with respect to the plane. The reflected surface lies in W .

We follow moving the plane on the left and reflecting the right part of S . Because S is compact, there would be a first time t such that the reflected surface touches the left part of S at some point p . We denote by Q_t this plane, S_t^- and S_t^+ each one of the left and right parts of S and let S_t^* be the reflection of S_t^+ with respect to Q_t . It can occur three possibilities (see figure 5.2):

1. If $p \in S_t^- \cap S_t^*$ is an interior point of both surfaces, the maximum principle says that $S_t^- = S_t^*$, that is, Q_t is a plane of symmetry of S .
2. If $p \in S_t^- \cap S_t^*$ is a point of $\partial S_t^- = \partial S_t^*$, we apply the maximum principle, in its boundary version.
3. If $p \in S_t^- \cap S_t^*$ and $p \in C$, then both surfaces are tangent since the surface makes constant angle with P along C . The maximum principle (in its interior or boundary version) says us that Q_t is a plane of symmetry again.




Figure 5.2: The Alexandrov reflection method.

Because the vector v is arbitrary, we have proved that given any vertical plane there exists other parallel plane which it is a symmetry plane of the surface. Since the surface is compact, we have showed that the surface is rotational with respect to a vertical straight-line.

Moreover, and from the proof, we deduce that the planes Q_t are planes of symmetry of C and the, C is a circle. Because the hyperbolic caps are the only CMC surfaces with circular boundary, our surface S must be a hyperbolic plane. $\square$

In Euclidean ambient there is not true the above result. It is necessary to add some extra hypothesis to assure that the surface is a planar domain or a spherical cap. For example that: i) the surface is a topological disc or; ii) the surface is embedded and lies in one side of the boundary plane.

5.2 The Dirichlet problem: the Euclidean case

The Dirichlet problem for the constant mean curvature equation is the following. Let $\Omega \subset \mathbb{R}^2$ be a domain, $\varphi \in C^0(\partial\Omega)$ and $H \in \mathbb{R}$. We ask for a solution $u = u(x, y)$ of the equation

$$\begin{cases} \operatorname{div} \frac{Du}{\sqrt{1 + |Du|^2}} = 2H & \text{on } \Omega, \\ u = \varphi & \text{in } \partial\Omega. \end{cases}$$

The first equation also writes as

$$(1 - u_y^2)u_{xx} - 2u_x u_y u_{xy} + (1 - u_x^2)u_{yy} = 2H(1 + u_x^2 + u_y^2)^{3/2}.$$

The surface $z = u(x, y)$ has mean curvature H with the upwards orientation

$$N = \frac{(-u_x, -u_y, 1)}{\sqrt{1 + |Du|^2}}.$$

Moreover, the boundary of S is the curve formed by the graph of φ .

From now, we suppose that $\varphi = 0$, that is, the boundary curve of the surface is included in a horizontal plane.

We use the continuity method to solve the problem. For this, we define the set

$$A = \left\{ t \in [0, H]; \text{ existe una solución del problema } \begin{array}{ll} \operatorname{div} \frac{Du}{\sqrt{1+|Du|^2}} = 2t & \text{en } \Omega \\ u = 0 & \text{en } \partial\Omega \end{array} \right\}.$$

This set is not empty since $0 \in A$: it suffices to take $u = 0$. If we prove that A is an open and closed set of $[0, H]$, then $A = [0, H]$. In particular, $H \in A$.

The study of the openness of A is a consequence of the Implicit Function Theorem. Exactly, for each immersion x and $u \in C_0^\infty(M)$ we define $x_t(p) = \exp_{x(p)}(tu(p)N(p))$. For t closed to 0, x_t is an immersion. Let H be the mean curvature. Moreover,

$$\frac{d}{dt}_{t=0} H(x_t(p)) := J(u)(p),$$

where J is the Jacobi operator given by

$$J = \Delta + 4H^2 - 2K.$$

Then $J\langle N, a \rangle = 0$ with $\langle N, a \rangle \geq 0$. This means that J is a Fredholm operator of index 0. The Implicit Function Theorem between Banach spaces asserts that the map $x \mapsto H(x)$ is invertible, that is, if t_0 is the mean curvature of a graph, there exists a neighborhood $(t_0 - \epsilon, t_0 + \epsilon)$ of t such that there exist graphs with mean curvature s with $s \in (t_0 - \epsilon, t_0 + \epsilon)$. In particular, t_0 is an interior point of A .

The proof of the closeness of A is a consequence to obtain a priori C^0 and C^1 estimates of a solution, that is, of $|u|$ and $|Du|$. When we say a priori we mean that given a solution u , we have to find a constant M , depending only on the initial conditions $(\Omega$ and $H)$ such that $|u|, |Du| < M$.

First, one can assure that there are C^0 estimates, that is, a priori bounds of $|u|$. We take the upwards orientation, that is, $\langle N, a \rangle > 0$, with $a = E_3$. Assume that $H > 0$: this means that the surface lies below the plane containing the boundary. By Theorem 4.3.1, we have

$$\Delta(H\langle x, a \rangle + \langle N, a \rangle) = -2(H^2 - K)\langle N, a \rangle \leq 0.$$

The maximum principle for elliptic equations asserts that

$$H\langle x, a \rangle + \langle N, a \rangle \geq \min_{\partial\Omega} (H\langle x, a \rangle + \langle N, a \rangle) = \min_{\partial\Omega} \langle N, a \rangle \geq 0.$$

By using that $x_3 := \langle x, a \rangle \leq 0$, we have proved that

$$Hx_3 \geq -\langle N, a \rangle \geq -1 \Rightarrow 0 \geq x_3 \geq \frac{-1}{H}.$$

This height bound depends only on H . Since $|u|$ measures the height of the graph of u , the above computation can read as

Proposition 5.2.1. *Let P be a plane and $H \in \mathbb{R}$. Given a graph S on P with constant mean curvature H and boundary included in P , the height of the graph is less than $1/|H|$.*

This means that, *given* a plane and a number H , the heights of the graphs with mean curvature H and boundary in the plane, have bounded heights.

We now study the C^1 estimates. First, we claim that it suffices to find bounds of $|Du|$ along $\partial\Omega$. Exactly, we are going to prove that

$$\max_{\Omega} |Du| = \max_{\partial\Omega} |Du|.$$

From the expression of the Laplacian of the function $\langle N, a \rangle$, we have

$$\Delta \langle N, a \rangle = -(4H^2 - 2K) \langle N, a \rangle \leq 0.$$

Then

$$\langle N, a \rangle \geq \min_{\partial\Omega} \langle N, a \rangle.$$

This means that

$$\frac{1}{\sqrt{1 + |Du|^2}} \geq \min_{\partial\Omega} \frac{1}{\sqrt{1 + |Du|^2}}.$$

Then we obtain the claim.

As a consequence, we want to estimate $|Du|$ *only* along $\partial\Omega$, that is, along the boundary curve of the surface. We write this in geometric terms. Let a be a unit vector orthogonal to the plane containing the boundary. Let ν be the interior conormal vector along ∂S . If $H > 0$, the surface lies below the plane. Moreover,

$$\langle \nu, a \rangle^2 + \langle N, a \rangle^2 = 1.$$

$$\langle \nu, a \rangle^2 = \frac{|Du|^2}{1 + |Du|^2} \Rightarrow |Du| = -\frac{\langle \nu, a \rangle}{\sqrt{1 - \langle \nu, a \rangle^2}}.$$

If $\delta = \min_{\partial\Omega} \langle \nu, a \rangle > -1$, then

$$\max_{\partial\Omega} |Du| = -\frac{\delta}{\sqrt{1 - \delta^2}}.$$

Let us see that $\langle \nu, a \rangle$ measures the slope of the surface with respect to the plane containing the boundary. Therefore, we have to find a maximum slope of the surface along the boundary.

An example of the above scheme is the following

Theorem 5.2.2. *Let Ω be a convex domain included in a plane P and let κ be the curvature of $\partial\Omega$. If $H \in \mathbb{R}$ satisfies $\kappa > H > 0$, then there exists a graph on Ω with mean curvature H and boundary $\partial\Omega$.*

Proof: Let r_1, r_0 be two positive numbers such that

$$\kappa > \frac{1}{r_1} > \frac{1}{r_0} > H.$$

Let K be a half-sphere of radius r_0 such that $K \subset \{z \leq 0\}$, $\partial K \subset P$ in such way that $\Omega \subset \Omega_1$, Ω_1 being the domain that bounds ∂K in P . This is possible since $r_0 > 1/\kappa$. We move upwards K a little in such way that in the new position, K' , the intersection of K' with P is a circle of radius r_1 and containing in its inside Ω . All this depends only on H and Ω . Let K^* be the spherical cap formed by the part of K' below P . This surface has the following properties:

1. It is a graph on P .
2. The circle ∂K^* has the property that can roll in such way it touches each one of the points of $\partial\Omega$.
3. The mean curvature of K^* is $1/r_0$, which is bigger than H .

Let G be a graph on Ω with mean curvature H and $\partial G = \partial\Omega$. We move downwards K^* until it does not touch G . Next, let us move upwards. By the maximum principle, K^* comes back to its original position without to contact with G . Moreover, we can move in horizontal direction in such way that it touches each point of ∂G with to be a tangent point. As a consequence, the slope of G is less than the one of K^* , obtaining the desired C^1 estimates.

□

5.3 The Dirichlet problem: the Lorentzian case

In Lorentz-Minkowski space, the corresponding Dirichlet problem is the following

$$\left\{ \begin{array}{ll} \operatorname{div} \frac{Du}{\sqrt{1-|Du|^2}} = 2H & \text{on } \Omega, \\ |Du| < 1 & \text{on } \partial\Omega. \\ u = 0 & \text{in } \partial\Omega. \end{array} \right.$$

The condition $|Du| < 1$ indicates that the graph is spacelike. The first difference with the Euclidean case is that the a priori estimates obtained in the Proposition 5.2.1 do not hold in our setting.

Proposition 5.3.1. *Let P be a plane and $H \in \mathbb{R}$. There exist graphs with constant mean curvature H , whose boundary lies in P with arbitrary heights.*

For this, it is suffices to consider hyperbolic caps $C(1/H; R)$, with R arbitrary and putting the boundaries of all them on the plane P .

Proposition 5.3.2. *Let Ω be a domain of a spacelike plane and $H \in \mathbb{R}$. Let $S = \operatorname{graph}(u)$ be a graph with mean curvature H and bounded by $\partial\Omega$. Then there exists a constant $c = c(H, \Omega)$ such that $|u| \leq c(H, \Omega)$.*

Proof: Without loss of generality, we assume that S is a graph on the plane $\{z = 0\}$ and that $\partial S \subset P$. Let $H > 0$ be the mean curvature. We know that $u \leq 0$. We take a hyperbolic cap $C(1/H; R)$, with the same mean curvature than S and orientation pointing upwards. We take R sufficiently big so that the boundary of $C(1/H; R)$ lies included in P and that Ω lies in the bounded domain determined by $\partial C(1/H; R)$.

By a translation, we move downwards $C(1/H; R)$ so that it does not touch S . Now, we move it up. Recall that the mean curvatures agree but not the boundary curves. Thus there is not a contact point unless that the hyperbolic cap returns its original position. This proves that the height of S is less than $C(1/H; R)$. $\square$

It is possible to obtain an estimate of the number $c(H, \Omega)$. Let $r_0 = \max_{\partial\Omega} \sqrt{x^2 + y^2}$. Then we take $R = 1 + r_0$ and the constant is c es $\sqrt{R^2 - 1/H^2} - 1/H$.

We now study the a priori C^1 estimates. Set $a = E_3 = (0, 0, 1)$. We know that

$$\Delta \langle N, a \rangle = (4H^2 + 2K) \langle N, a \rangle.$$

As the surface is a spacelike graph with the upwards orientation, then $\langle N, a \rangle \leq -1$. On the other hand, $4H^2 + 2K \geq 2H^2 \geq 0$. Then $\Delta \langle N, a \rangle \leq 0$. This implies that

$$\langle N, a \rangle \geq \min_{\partial\Omega} \langle N, a \rangle, \quad \text{con } \langle N, a \rangle = -\frac{1}{\sqrt{1 - |Du|^2}}.$$

This means that

$$\max_{\Omega} |Du| = \max_{\partial\Omega} |Du| < 1$$

In this moment we can continue in Euclidean language: in order to find a priori C^1 estimates, we have to find an upper bound $c = c(H, \Omega)$ of the slope of the surface with $c(H, \Omega) < \sqrt{2}/2$.

We have the next

Theorem 5.3.3. *If Ω is a bounded, convex domain, then the Dirichlet problem has a solution (for any H).*

Proof: The proof is analogous to Theorem 5.2.2, but working with hyperbolic caps. Let $H \in \mathbb{R}$ and r_0 the number defined in that theorem. Let $S = C(1/H; R)$ be a hyperbolic cap with $\partial S \subset P$ and $R > 1 + r_0$. Let $D \subset P$ be the domain that determines ∂S . Let R be sufficiently big so that $D \supset \overline{\Omega}$ and the circle ∂S can roll touching each point of $\partial\Omega$. This radius depends only on H and Ω .

We have proved that S lies in the domain bounded by $S \cup D$. By horizontal translations, we can carry S in such way that ∂S touches each point of $\partial\Omega$ without to be tangent (by the maximum principle). Therefore, the slope of G is less than the one of $C(1/H; R)$, which it is less than $\sqrt{2}/2$. $\square$

5.4 Exercises

1. Show that it is not possible to obtain an estimate of C^0 type depending only on H with the same reasoning as in the Proposition 5.2.1.

2. Let S_1 and S_2 be two graphs in Euclidean space $\mathbf{E}^3$ with constant mean curvature H_1 and H_2 respectively (with upwards orientations in both surfaces). Let $S_i = \text{graph}(u_i)$ and assume that $\partial S_1 = \partial S_2$. Then if $H_1 > H_2$, then $u_1 < u_2$ in Ω . Find the corresponding Lorentzian version.

Chapter 6

Lorentzian Riemann examples

In this chapter we are going to show the spacelike CMC surfaces of $\mathbf{E}_1^3$ foliated by circles. As a particular case of this kind of surfaces are the surfaces of revolution. In contrast to what occurs in Euclidean space, in Minkowski space we have three types of rotational surfaces depending on the causal character of the axis of revolution.

6.1 Introduction to the problem

A way to obtain examples of surfaces with some condition on the mean curvature is to impose a shape to the surface. In this chapter we consider that the surface is rotational, that is, it is invariant by a group of rotations that leave pointwise fixed a straight-line of the space. This implies that the mean curvature equation is now an ordinary differential equation. Under appropriate initial conditions, there exists a such surface.

In Euclidean space, and if $H = 0$ (minimal surface), planes and catenoids are the only rotational surfaces with $H = 0$. The parametrization of the catenoid is (up isometries and homotheties)

$$X(s, \theta) = (\cosh(s) \cos(\theta), \cosh(s) \sin(\theta), s),$$

where the z -axis is the axis of revolution. See figure 6.1.

If one asks for minimal surfaces formed by a uniparametric family of circles in parallel planes, then there are other surfaces: they are the Riemann examples¹. These surfaces are foliated by circles in parallel planes and in a discrete set of heights, the surface is asymptotic to horizontal planes. See figure 6.1

Motivated by the Riemann examples, we generalize the notion of rotational surface.

Definition 6.1.1. *A surface in Euclidean space is called cyclic if it is formed by a uniparametric family of circles, that is, it is possible to locally parametrize as*

$$X(t, s) = c(t) + r(t)(\cos(s)e_1(t) + \sin(s)e_2(t)),$$

¹All our pictures have been made by using Mathematica. The equations that describe such surfaces are analogous to the Lorentzian case and for this reason, we left to the reader the work to find them




Figure 6.1: The catenoid and one Riemann example of $\mathbf{E}^3$.

where $c(t) \in \mathbb{R}^3$, $r(t) > 0$ y $\{e_1(t), e_2(t)\}$ are orthonormal vectors (all functions that appear are differentiable).

One can prove that if a minimal surface is foliated by circles, then these circles lie in parallel planes. Assume that H is constant, but $H \neq 0$. Then

Theorem 6.1.2. *The only cyclic surfaces of $\mathbf{E}^3$ with constant mean curvature $H \neq 0$ are the surfaces of revolution.*

This means that "there are not Riemann examples when $H \neq 0$ ". Let us see that the planes of the foliation are parallel, but that any intersection of a sphere with a family of (non-parallel) planes are circles.

We now consider the Minkowski space $\mathbf{E}_1^3$. We pose the corresponding

Problem. Find all cyclic CMC spacelike surfaces in Lorentz-Minkowski space.

See the paper [3] as a short survey about this problem. The same question can be posed for timelike surfaces. As we have pointed out at the beginning of this chapter, the difference with the Euclidean case is that the axis of revolution can be of three causal types. In Chapter 1 we have studied the group of isometries that leave pointwise fixed a straight-line L (a group of boosts). Moreover, we have studied the orbit of a point that does not lie in the axis of revolution. With respect to an appropriate base $\{e_1, e_2, e_3\}$, we have:

1. if L is timelike, $L = \langle e_3 \rangle$, the group of boosts that leave fixed L is $\{R_\theta; \theta \in \mathbb{R}\}$

$$R_\theta = \begin{pmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

The circles write as

$$\alpha(s) = c + r(\cos(s) e_1 + \sin(s) e_2), \quad (6.1)$$

with $r \neq 0$, $c \notin L$.

2. If $L = \langle e_1 \rangle$ is spacelike, the group is

$$R_\theta = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cosh \theta & \sinh \theta \\ 0 & \sinh \theta & \cosh \theta \end{pmatrix}.$$

The circles are

$$\alpha(s) = c + r(\sinh s \, e_2 + \cosh s \, e_3), \quad (6.2)$$

with $r > 0$ and $c \notin L$.

3. If L is lightlike, we consider $L = \langle e_2 + e_3 \rangle$, with e_3 a timelike vector. The group is

$$R_\theta = \begin{pmatrix} 1 & \theta & -\theta \\ -\theta & 1 - \frac{\theta^2}{2} & \frac{\theta^2}{2} \\ -\theta & -\frac{\theta^2}{2} & 1 + \frac{\theta^2}{2} \end{pmatrix}$$

and the circles are

$$\alpha(s) = c + s e_1 + \frac{r s^2}{2} (e_2 + e_3), \quad (6.3)$$

After an isometry of the ambient space $\mathbf{E}_1^3$, and if we take the usual base, one has

1. if $L = \langle E_3 \rangle$, the circles are Euclidean circles in horizontal planes.
2. if $L = \langle E_1 \rangle$, the circles are hyperbolas in parallel planes to $\langle E_2, E_3 \rangle$.
3. if $L = \langle E_2 + E_3 \rangle$, the circles are parabolas in parallel planes to $\langle E_1, E_2 + E_3 \rangle$.

Definition 6.1.3. A rotational surface of $\mathbf{E}_1^3$ is a surface invariant by a group of rotations that leaves pointwise fixed a straight-line L .

Our work plan is the following:

1. Study cyclic CMC surfaces where the planes of the foliation are parallel. Show that the surface is rotational or $H = 0$.
2. Show that cyclic CMC surfaces are of the above kind.
3. Study the rotational CMC surfaces.

6.2 The planes of the foliation are parallel

The main result that we obtain is the following

Theorem 6.2.1. Let S be a spacelike surface foliated by circles in parallel planes. If the mean curvature H is constant, then

1. If $H \neq 0$, the surface is rotational.

2. If $H = 0$, the surface is rotational or it belongs to the family of Riemann examples.

By a "Riemann example" we mean a non-rotational cyclic surface with mean curvature $H = 0$ and the circles lie in parallel planes. At certain heights, the surface has a singularity or it is asymptotic to a plane.

The proof of Theorem must distinguish the kinds of circles. In order to show how is the reasoning, we only do the case that the foliation planes are spacelike (and the circles are Euclidean circles).

We parametrize the surface as

$$X(u, v) = (a + r \cos(v), b + r \sin(v), u),$$

where $a, b, r > 0$ are smooth functions of u . The curve $u \mapsto (a(u), b(u), u)$ describes the centers of the circles. The fact that the surface is rotational is equivalent that the functions a and b are constant.

We compute the mean curvature. From the Chapter 3, we know that H writes as

$$2HW^{3/2} = E \det(X_u, X_v, X_{vv}) - 2F \det(X_u, X_v, X_{uv}) + G \det(X_u, X_v, X_{uu}) := P, \quad (6.4)$$

where $W = EG - F^2$. We distinguish two cases:

1. Case $H \neq 0$. After a homothety, we assume that $H = 1/2$. Squaring equation (6.4), we obtain $W^3 - P^2 = 0$. After a long computation², this equation writes as a polynomial of type

$$\sum_{n=1}^6 A_n(u) \cos(nv) + B_n(u) \sin(nv) = 0. \quad (6.5)$$

As the functions $\cos(nv)$ and $\sin(nv)$ are independent linear, the functions A_n and B_n must vanish. From $A_5 = B_5 = 0$ we obtain

$$a'^4 - 10a'^2b'^2 + 5b'^4 = 0, \quad 5a'^4 - 10a'^2b'^2 + b'^4 = 0.$$

Hence we deduce that $a' = b' = 0$, that is, a and b are constant. This shows that the surface is rotational. In such case, and taking $a = b = 0$, the mean curvature satisfies the equation

$$H = \frac{-1 + r'^2 - rr''}{2r(-1 + r'^2)^{3/2}}.$$


A picture of such surface for $H = 1$ appears in figure 6.2.

2. Case $H = 0$. Then $P = 0$ gives a polynomial as (6.5), but only until $n = 1$. The coefficients A_1 and B_1 give

$$2a'r' - ra'' = 0, \quad 2b'r' - rb'' = 0.$$

A first integration concludes that there exist constants c and d such that $a' = cr^2$ y $b' = dr^2$.

²We use Mathematica.

Figure 6.2: A rotational surface with $H = 1$.

- (a) If $c = d = 0$, the surface is rotational. The equation that describes this surface, named Lorentzian catenoid, is

$$-1 + r'^2 - rr'' = 0.$$

This equation can integrate and, after homotheties, the solution is $r(s) = \sinh(s)$. Therefore, the Lorentzian catenoid parametrizes as

$$X(s, \theta) = (\sinh(s) \cos(\theta), \sinh(s) \sin(\theta), s)$$

and its picture can see in the figure 6.3.




Figure 6.3: The Lorentzian catenoid.

- (b) If $cd \neq 0$, the equation that satisfies r is

$$-1 + (c^2 + d^2)r^4 + r'^2 - rr'' = 0.$$

The solutions of this equation give the Riemann examples³. See figure 6.4. This surface appeared in the author's work: R. López, Constant mean curvature hypersurfaces foliated by spheres, *Differential Geometry and its Applications*, 11 (1999), no. 3, 245–256.




Figure 6.4: Riemann examples in $\mathbf{E}_1^3$. These surfaces are foliated by Euclidean circles in parallel horizontal planes.

Finally, we remark that a similar work can do for surfaces foliated by hyperbolas and parabolas. In the case that the surface is foliated by hyperbolas, we have the corresponding Lorentzian catenoid and Riemann examples. See figure 6.5.




Figure 6.5: A rotational surface with $H = 0$ and foliated by hyperbolas. On the right, a Riemann example foliated by hyperbolas.

6.3 The planes of the foliation are not parallel

We now suppose that S is a spacelike surface foliated by circles in non-parallel planes. Let H be the value of the (constant) mean curvature. The reasonings for $H = 0$ and $H \neq 0$ are similar. If $H = 0$, we conclude that it is not possible this situation; if $H \neq 0$, we obtain that the surface must be a hyperbolic plane. We do the reasoning for surfaces foliated by spacelike planes.

³Let us see that for initial conditions, for example, at $s = 0$, the value of W must be positive, since the surface is spacelike. In our case, this means $(a' + r')^2 > 1$

Let $\Pi = \Pi(u)$ be the planes of the foliation parametrized by the parameter u . Suppose that the planes are not parallel. We consider the curve $\Gamma = \Gamma(u)$ orthogonal to each plane $\Pi(u)$, where u denotes the length-arc parameter of Γ . Let κ be the curvature of Γ . Because Γ is not a straight-line, $\kappa \neq 0$ (our reasoning is local). Let $\mathbf{T}$ be the tangent vector of Γ . Then $\mathbf{T}$ is timelike and the Frenet equations of Γ are

$$\begin{aligned}\mathbf{T}' &= \kappa \mathbf{N} \\ \mathbf{N}' &= \kappa \mathbf{T} + \sigma \mathbf{B} \\ \mathbf{B}' &= -\sigma \mathbf{N}\end{aligned}$$

The surface can parametrize as

$$X(u, v) = \mathbf{c} + r(\cos v \mathbf{N} + \sin v \mathbf{B}),$$

where $r = r(u) > 0$ and $\mathbf{c}(u)$ are differentiable functions. The curve $\mathbf{c} = \mathbf{c}(u)$ is the curve that describes the centers of circles. We write

$$\mathbf{c}' = \alpha \mathbf{T} + \beta \mathbf{N} + \gamma \mathbf{B}, \quad (6.6)$$

where α, β and γ are smooth functions on u . We compute again W and P and we distinguish the cases $H = 0$ and $H \neq 0$. Assume that $H \neq 0$. After a homothety, we assume that $H = 1/2$ and that $P^2 - W^3 = 0$. The expressions of P and W are now a polynomial on $\cos nv, \sin nv$. Exactly,

$$P^2 - W^3 = \sum_{i=0}^{12} \left[A_i(u) \cos(nv) + B_i(u) \sin(nv) \right] = 0.$$

The computation of the coefficients is very hard, being necessary the use of a symbolic programme as Mathematica.

By the amount of computations and cases, we are not going to do the complete reasoning. We point out that the contradiction arrives when one conclude that $\kappa = 0$ or that $W = EG - F^2 = 0$. In order to show how one obtain hyperbolic planes, we explicit what happens in this situation. In general, the coefficient B_{12} is

$$B_{12} = \frac{r^{12} \tau^4}{2048}.$$

Thus $\tau = 0$, that is, the curve of the circle centers is a planar curve.

From now, the next coefficients vanish until that $n = 6$. From $B_6 = 0$, we have some possibilities. We analyze one of them and that corresponds with

$$\gamma = 0 \quad \text{y} \quad \beta^2 = \kappa^2(4 + r^2).$$

Now B_5 yields

$$B_5 = \kappa^5 r^7 \left(\alpha - \frac{rr'}{\sqrt{4 + r^2}} \right) = 0.$$

Then $\alpha = rr' / \sqrt{4 + r^2}$. Hence, the equation $W^3 - P^2 = 0$ is trivial. With the values of α, β and γ , the derivative of the center curve $\mathbf{c}$ is:

$$\mathbf{c}' = \frac{rr'}{\sqrt{4 + r^2}} \mathbf{T} + \kappa \sqrt{4 + r^2} \mathbf{N} = (\sqrt{4 + r^2} \mathbf{T})'.$$

This expression can integrate, obtaining that there is $c_0 \in \mathbb{R}^3$, such that

$$\mathbf{c} = c_0 + \sqrt{4 + r^2} \mathbf{T}.$$

The expression of the parametrization $X(u, v)$ of the surface writes now as

$$X(u, v) = c_0 + \sqrt{4 + r^2} \mathbf{T} + r \cos(v) \mathbf{N} + r \sin(v) \mathbf{B}.$$

Hence we deduce

$$\langle X - c_0, X - c_0 \rangle = -4,$$

that is, it is a hyperbolic plane with mean curvature $H = 1/2$.

6.4 Exercises

1. Find the equations that describe the Riemann examples in Euclidean space.
2. Find the equations that describe the rotational (spacelike and timelike) surfaces with constant mean curvature.

Bibliography

1. M. P. do Carmo, Differential Geometry of Curves and Surfaces, Prentice Hall, 1976.
2. D. Gilbarg, N. Trudinger, Elliptic Partial Differential Equations of Second Order, Springer-Verlag, 1983.
3. R. López, Cyclic hypersurfaces of constant curvature, Advanced Studies in Pure Mathematics, 34, 2002, Minimal Surfaces, Geometric Analysis and Symplectic Geometry, 185–199.
4. B.O'Neill, SemiRiemannian geometry with application to general relativity, Academic. Press, New York, 1983.
5. T. Weinstein, An Introduction to Lorentz Surfaces, Walter de Gruyter, 1996.

Index

- a priori estimates, 51
- Alexandrov reflection method, 49
- arc-length, 19
- area functional, 40

- Bertrand curve, 27
- binormal vector, 21, 22
- boost, 11, 24

- catenoid, 57
- Cauchy-Schwarz inequality, 7
- circle, 13, 25
- comparison principle, 42
- continuity method, 51
- curvature, 21
- curvature of a curve, 21
- cyclic surface, 57

- De Sitter space, 31, 37
- Dirichlet problem, 50, 53

- Frenet equations, 21, 22

- Gauss curvature, 34
- Gauss formula, 33
- Gauss map, 32

- helix, 26
- hyperbola, 13, 25
- hyperbolic angle, 8
- hyperbolic cap, 48
- hyperbolic plane, 30, 37

- isometry, 10

- Jacobi operator, 51

- Laplacian, 44
- lightlike
 - curve, 17
- immersion, 30
 - subspace, 2
 - vector, 2
- lightlike cone, 2, 31
- linear elliptic equation, 44
- Lorentz-Minkowski space, 1
- Lorentzian catenoid, 61, 62
- Lorentzian metric, 1

- mate curve, 27
- maximum principle, 43
- mean curvature, 33
- modulus, 6

- normal vector, 21, 22

- orientation, 9
- ortocrone group, 10

- parabola, 13, 25
- principal curvatures, 33
- pseudo parametrization, 20

- quasilinear elliptic equation, 44

- Riemann examples, 60, 62

- second fundamental form, 33
- spacelike
 - curve, 17
 - immersion, 30
 - subspace, 2
 - vector, 2
- special Lorentz group, 10
- special Lorentz ortocrone group, 10

- tangent point, 42
- tangent vector, 21
- timelike
 - curve, 17

- immersion, 30
 - subspace, 2
 - vector, 2
- timelike cone, 6
- timelike orientation, 9
- torsion, 21, 22
- totally umbilical, 36
- trihedron Frenet, 21, 22
- umbilical point, 36
- variation of a immersion, 39
- variational vector field, 39
- volume functional, 40
- Weingarten endomorphism, 33